

A Comparative study of storage stability in virgin coconut oil and extra virgin Olive oil upon thermal treatment

*Henna Lu, F. S. and Tan, P.P.

Department of Food Science and Nutrition, Faculty of Applied Sciences, UCSI University, 56000 Kuala Lumpur, Malaysia

Abstract: This aim of this research is to investigate thermal stability of virgin coconut oil, (VCO) which was heated at 190°C upon 40 days storage as compared to extra virgin olive oil (EVOO). The changes in fatty acids composition through (GC), Fourier Transform Infrared (FTIR) spectra, iodine value (IV) and total phenolic content were determined throughout the period of study. Results from GC showed that there was significant changes ($P<0.05$) in the percentage of linolenic acid (from 0.74% to 0.54%) and palmitic acid (from 12.31% to 13.44%) in the EVOO upon storage. Higher stability was found in VCO as it contains higher saturated fatty acids. FTIR spectra showed both oils had undergone oxidation and produced hydroperoxides upon 40 days storage but with less oxidation in VCO. Meanwhile, IV and total phenolic content decreased significantly ($P<0.05$) in both oils with percentage reduction of IV in VCO greater than that in EVOO (70.60% vs. 11.90%), contrary to percentage loss in total phenolic content (50.77% vs. 68.19%). As a conclusion, VCO can be considered as good frying oil as it has relatively high oxidative stability as EVOO.

Keywords: Virgin coconut oil, extra virgin olive oil, oil stability, FTIR, IV, GC

Introduction

Frying is a common unit operation used in the preparation and manufacture of food. It is the process of cooking and drying through contact with hot oil and simultaneous heat along with mass transfer is applied (Fellows, 2003). During frying, cooking oil which acts as heat transfer media is exposed to elevated temperature constantly in the presence of air and moisture. Under such conditions, some complex reactions like polymerization, oxidation and hydrolysis take place. These reactions influence many qualities of the finished product in terms of flavor, color, texture, shelf life and nutritional aspects (Boskou, 2003). Thus, the selection of oil that is as stable to reactions above as possible is very crucial. Most vegetable oils such as safflower oil, sunflower oil, soybean

oil and corn oil consist of high percentage of polyunsaturated fatty acids which causes them to be more susceptible to deterioration in quality and less suitable for frying. Monounsaturated fatty acids, on the other hand, more stable and oxidize less readily than polyunsaturated ones (Mackay, 2000). Extra virgin olive oil (EVOO) is an excellent example of oil that is rich in oleic acid, a monounsaturated fatty acid. Its relatively high oxidative stability is also contributed by its high tocopherol and phenolic content (Andrikopoulos *et al.*, 2002; Gomez-Alonso *et al.*, 2003). EVOO is the product of mechanical extraction from fresh ripe fruits of olive tree with no heat involved and no chemicals such as synthetic antioxidants added to it. Similar to EVOO, virgin coconut oil (VCO) is produced naturally from fresh, mature kernel of coconut. Since minimal heat and process is involved, VCO retains its

*Corresponding author.

Email: fslu@ucsi.edu.my

Tel: +603-91018880; Fax: +603-91023606

richness of antioxidant content too. VCO is gaining the popularity in scientific field and among the consumer.

Recent studies shown that consumption of VCO has a beneficial effect in lowering harmful lipid components compared to refined, bleached and deodorized (RBD) coconut oil. VCO is not only able to reduce the level of total cholesterol, triglycerides, phospholipids, low-density lipoprotein (LDL) and very-low-density lipoprotein (VLDL) cholesterol, but also capable to increase the concentration of high density lipoprotein (HDL), which is good cholesterol (Nevin and Rajamohan, 2004). Another animal study done by Nevin and Rajamohan (2006) showed that consumption of VCO which contains more unsaponifiable components like vitamin E and polyphenols than RBD coconut oil, resulted in an increased level of antioxidant enzymes and a reduced lipid peroxide content in both in vitro and in vivo conditions. Although VCO is classified under vegetable oils just as EVOO, it consists of high percentage of saturated fatty acids which are usually present abundantly in oils made from palm as well as animal sources. Along with high antioxidant content, the high degree of saturation in fatty acids of VCO may further enhance its oxidative stability. To date, studies done on VCO were focused on its beneficial health effects (Nevin and Rajamohan, 2004; 2006) and sensory evaluation (Villarino *et al.*, 2005), no literature on the stability of VCO. The objective of present work is to determine the suitability of VCO as frying oil by investigating its thermal stability upon storage through its fatty acids composition, total phenolic content and Iodine Value which show oxidative level of the oil.

Materials and Methods

Materials

Virgin coconut oil, VCO (Country Farm Organic, Malaysia), extra virgin olive oil, EVOO (Borges, Spain) and French fries used for frying were purchased from local supermarket.

Heat treatment and storage studies

The oils were heated to speed up the onset of degradative reactions such as hydrolysis, oxidation and polymerization in the oils used in this study. Frying was conducted in a Philips deep fryer (Cucina DH 6151, China) equipped with thermostat. First, 500 ml of oil sample was preheated to a temperature of 190°C. After 15 min, 30 g of raw French fries was added. The French fries were fried for 5 min and then removed. Following this procedure, the oil temperature was allowed to return to 190°C within 5 min before subsequent frying. Each of the oil samples completed six frying operations of French fries, equivalent to a period of 75 min. The oil was allowed to cool down to room temperature before analysis was carried out. Oil samples after completion of 6 frying operations were filled into test tubes with screw cap. The entry of oxygen was further prevented by wrapping the tubes with parafilm. Additionally, the tubes were covered with a layer of aluminum foil to exclude the light completely and then stored at room temperature. Samples were taken out for analysis every tenth day for a total storage period of 40 days.

Determination of FAMEs composition by Gas Chromatography

Perkin Elmer Gas Chromatograph equipped with Flame Ionization Detector (Elmer Instruments Autosystem XL) was used to determine the FAMEs composition of the tested oils. The column used was fused silica capillary (SPTM 2380, Supelco; 30 m x 0.25 mm x 0.25 µm film thickness).

Flow rates of compressed air and hydrogen were at 450ml/min and 25 ml/min respectively while the nitrogen gas was at the pressure of 180 psi. The split ratio used was 50:1. Initial oven temperature was 140°C, held for 5 min then programmed to 240°C at 4°C per min, and finally held for 10 min at this temperature. The temperatures of injector and FID were both set at 250°C. Total runtime was 40 min. The oil was converted to fatty acid methyl esters (FAMEs) according to the modified method described by Qian (2003). First, 0.5 g of oil was dissolved in 25 ml n-hexane. Then, 5mL of the mixture was transferred into a test tube and mixed with 250 μ l 0.5M sodium methoxide in methanol. The mixture was then shaken vigorously in a vortex mixer for 60 seconds. Five milliliters of saturated sodium chloride solution was then added and the mixture was again shaken vigorously for 15 seconds. After 10 min, 3 ml of hexane layer containing FAME (the top layer) was transferred into a vial followed by addition of small amount of anhydrous sodium sulfate. Finally, approximately 1 μ l of sample was injected manually into the gas chromatograph. The above procedures were performed in duplicate in fresh oil samples as well as after cooling down (0 day) and in the end of experimentation (stored for 40 days) in heated oil samples. The chromatographic data was recorded and analyzed in a personal computer (Optiplex GX 160, Dell) using TotalChrom Navigator Software, Perkin Elmer, USA. Individual fatty acids were identified by comparing their retention time with standard FAME mix GLC-10 and Standard Supelco-37 components FAMEs. The percentage of each component was calculated by dividing the individual peak area with total peak area.

Collection of FTIR spectra

Potassium bromide (KBr) powder was ground by using the grinder specialized for FTIR spectrometer. Then the KBr powder was heated in an oven at 100°C for 30 min to reduce the moisture content. By using the presser specialized for FTIR spectrometer, the KBr powder was pressed strongly to form a transparent KBr disk. The KBr disk was then placed in the standard holder specialized for FTIR spectrometer. A background spectrum was first obtained by analyzing the KBr disk. Next, the KBr disk was taken out and a small drop of oil was placed on KBr disk to form a thin capillary film. The disk was placed in the holder again for analysis. FTIR spectra of the oil samples were collected through FTIR spectrometer Thermo Nicolet-Avatar 320. The infrared spectra were recorded in the range 4000 cm^{-1} to 400 cm^{-1} . All data were recorded and integrated in a personal computer operating under Nicolet Omnic software version 6.

Iodine value

Iodine value was determined the AOAC Official Methods of Analysis (1984), with minor modifications. Approximately 0.3 g of sample was weighed into a 250 ml conical flask followed by addition of 10 ml of carbon tetrachloride. Twenty five milliliters of Wijs reagent was added into the flask. The contents were mixed thoroughly by swirling and stored in a dark place at room temperature. At the end of 30 min, 20 ml of 10% potassium iodide solution and 100 ml of distilled water were added to the sample solution. The mixture was titrated immediately with standard 0.25M sodium thiosulfate solution until the yellow color almost disappears. Three milliliters of starch indicator solution was added and titration was carried on dropwise with vigorous swirling until disappearance of the blue starch-iodine color. The blank was prepared by adding 10 ml of carbon tetrachloride into an additional flask without the oil. The blank

was then titrated in the same manner as described previously. The iodine value was then calculated by using the formula as described by Xu *et al.* (2007). IV test was performed in duplicate for each of the oil samples.

Total phenolic content

The phenolic content of oil was extracted according to the method described by Rotondi *et al.* (2004) with some modifications. Approximately 15 g of oil was weighed into a 50 ml Falcon tube. Ten milliliters of n-hexane was mixed with the oil. The mixture was extracted with 10 ml of methanol: water (60:40). The mixtures were shaken for 5 min and then centrifuged at 5500 rpm for 5 min. The hydroalcoholic phase was collected and the hexane phase was re-extracted twice with 10 ml of methanol: water (60:40) each time. The combined hydroalcoholic fractions from three extractions were subjected to final washing with 10 ml of n-hexane to remove residual oil in a separatory funnel. The excess solvent was evaporated under vacuum at 40°C until dryness in a rotary evaporator. The residue was reconstituted in 2 ml methanol: water (60:40) for VCO sample and 20 ml methanol: water (60:40) for EVOO sample. The difference in amount of methanol:water (60:40) used was due to large difference in amount of phenolic compounds present in the samples.

The total phenolic content was determined by the Folin-Ciocalteu reagent assay (Lim *et al.* 2007). First, 0.5 ml of the extract obtained was mixed with 1.5 ml of Folin-Ciocalteu reagent previously diluted with distilled water (1:10). After standing at room temperature for 3 min, 1.2 ml of 15% sodium carbonate solution was added. The mixture was placed in dark room for 60 min. After that, absorbance was measured at 765 nm against the blank using a spectrophotometer (Secomam UVi light

XTD). The calibration curve was obtained by repeating the above procedures with known concentrations of gallic acid solutions. The results were expressed as milligrams of gallic acid equivalents (GAE) per 100 gram of oil (mg GAE per 100 g of oil). Since the assay quantifies all phenolic compounds, the selection of gallic acid as a standard is based on the availability of a stable and pure compound. In addition, gallic acid is cheaper than other options. Analyses were performed in duplicate for each of the extract.

Statistical analysis

Since all analyses were done in duplicate, the presented results are the mean of the obtained values. Minitab 15 statistical software was used for statistical analyses of fatty acids profile, IV and total phenolic content. Analysis of variance (ANOVA) and Fisher LSD mean comparison test were applied. Significant differences were detected at ($P \leq 0.05$).

Results and Discussion

Changes in fatty acids composition upon storage

Vegetable oils undergo changes in terms of chemical and physical properties when they interact with the food or atmosphere. Types of reaction that are known to lead to degradation of vegetable oils include polymerization, oxidation and hydrolysis. The changes in fatty acids composition of EVOO and VCO are shown in Table 1. In EVOO, the fatty acid composition suffered slight changes upon heating and storage. There was a significant decrease ($P < 0.05$) in percentage of linolenic acid after heat treatment while that of palmitic acid increased significantly ($P < 0.05$) with storage time. This trend was due to lipid oxidation in the oil and was in good

Table 1. Changes in fatty acids composition in VCO and EVOO

Retention Time (minute)	Fatty acids	VCO* (%)			EVOO* (%)		
		Before heat treatment (fresh)	After heat treatment (day 0)	After heat treatment (day 40)	Before heat treatment (fresh)	After heat treatment (day 0)	After heat treatment (day 40)
1.41	Caprylic acid (C8:0)	9.00 ± 0.76 ^a	10.18 ± 1.67 ^a	10.55 ± 1.29 ^a			
2.09	Capric acid (C10:0)	5.66 ± 1.35 ^a	7.05 ± 0.74 ^a	6.23 ± 0.85 ^a			
3.81	Lauric acid (C12:0)	51.31 ± 2.94 ^a	59.79 ± 1.99 ^a	48.63 ± 2.21 ^a			
7.44	Myristic acid (C14:0)	18.50 ± 0.16 ^a	16.69 ± 2.54 ^a	17.54 ± 0.38 ^a			
11.99	Palmitic acid (C16:0)	7.76 ± 0.34 ^a	8.12 ± 1.01 ^a	9.02 ± 0.01 ^a	12.31 ± 0.28 ^a	13.34 ± 0.24 ^{ab}	13.44 ± 0.20 ^b
12.55	Palmitoleic acid (C16:1)				0.85 ± 0.02 ^a	0.89 ± 0.06 ^a	0.86 ± 0.01 ^a
16.60	Stearic acid (C18:0)	2.71 ± 0.24 ^a	2.64 ± 0.41 ^a	3.03 ± 0.14 ^a	3.61 ± 0.18 ^a	3.75 ± 0.21 ^a	4.30 ± 0.06 ^a
17.00	Oleic acid (C18:1)	4.31 ± 0.37 ^a	4.09 ± 0.43 ^a	4.69 ± 0.09 ^a	75.21 ± 0.70 ^a	74.63 ± 0.72 ^a	74.12 ± 0.26 ^a
17.97	Linoleic acid (C18:2)	0.77 ± 0.15 ^a	0.42 ± 0.00 ^a	0.31 ± 0.09 ^a	7.30 ± 0.24 ^a	6.78 ± 0.12 ^a	6.76 ± 0.06 ^a
19.37	Linolenic acid (C18:3)				0.74 ± 0.02 ^a	0.58 ± 0.02 ^b	0.54 ± 0.04 ^b
	SFA	94.93	95.49	95.00	15.92	17.09	17.74
	PUFA	0.77	0.42	0.30	8.03	7.39	7.29
	MUFA	4.30	4.09	4.70	76.05	75.52	74.95

*Each value from the table represents the mean ± standard deviation of duplicate results.

Means within each row for each oil with different superscripts are significantly different (P<0.05).





Figure 1.Chromatograms of EVOO (a) Fresh; (b) After heat treatment at day 0; and (c) After heat treatment at day 40.

Table 2. FTIR absorbance bands and their characteristic functional groups

Number	Wave numbers (cm^{-1})	Characteristic group and mode of vibration
1	3470	Overtone of $-\text{C=O}$ (ester)
2	3006	Stretching of $=\text{C}-\text{H}$ (cis)
3	2925	Asymmetric stretching of $-\text{CH}$ (CH_2)
4	2854	Symmetric stretching of $-\text{CH}$ (CH_2)
5	1746	Stretching of $-\text{C=O}$ (ester)
6	1655	Stretching of $-\text{C-C-}$ (cis)
7	1464	Bending (scissoring) of $-\text{C-H}$ (CH_2 and CH_3)
8	1377	Symmetric bending of $-\text{C-H}$ (CH_3)
9	1237	Stretching, bending of $-\text{C-O}$, $-\text{CH}_2-$
10	1163	Stretching, bending of $-\text{C-O}$, $-\text{CH}_2-$
11	1099	Stretching of $-\text{C-O}$
12	723	Bending (rocking) of $-(\text{CH}_2)_n-$, HC=CH- (cis)

Source: Guillén and Cabo (1997)

**Figure 3.** FTIR spectra of fresh extra virgin olive oil (EVOO) and virgin coconut oil (VCO).

agreement with that reported by Warner and Mounts (1993) and Chung *et al.* (2006). The significant decrease in percentage of highly unsaturated fatty acids (in this case, linolenic acid) is due to the fact that they are more susceptible to oxidation than other fatty acids. The other fatty acids experienced no significant changes in terms of their percentage. The changes in percentage of individual fatty acids in EVOO do not only account for chemical reactions that occur during frying and storage, but also due to the leaching of the oils from French fries to EVOO during frying. On the other hand, the fatty acids composition of VCO remained

fairly constant throughout the experiment. In VCO, the relative content of unsaturated fatty acids was small to produce any detectable changes caused by high temperature and storage, and thus did not alter the proportion significantly. High percentage of saturated fatty acids in VCO caused this oil to have higher stability as compared to EVOO which has lower percentage of saturated fatty acids but higher unsaturated fatty acids. However, it should be noted that a constant fatty acid composition presented in terms of percentage does not necessarily indicate that the concentration of particular fatty acids

remain unchanged. The chromatograms of EVOO and VCO at different stages during the study are shown in Figure 1 and 2 respectively. There were noticeable decreases in the peak height and peak area of each component in both oil samples, which may indicates that the oils were degrading and broken down. This proved that the concentration of each fatty acid was actually decreasing despite the fact the exact reduction was not calculated quantitatively due to certain limitation in this preliminary study. The degradations resulted from oxidation and other reactions such as polymerization that take place during heat treatment and storage.

FTIR spectra for fresh oils

FTIR shows a few advantages over classical parameters in the measurements of oil stability. With only a small drop of oil and a couple of minutes to prepare the sample, the infrared spectrum collected can give information on different functional groups present. Also, minimal errors in the determination of frequency and absorbance can be expected since it is fully computerized (Guillén and Cabo, 2004). In the characterization of vegetable oils, the bands of interest are absorption peaks at 3600-2800 and 1800-700 cm^{-1} . The band assignments and their respective mode of vibration are shown in Table 2 and Figure 3. As clearly indicated in Figure 3, there were visible peaks near 3006 and 1655 cm^{-1} in spectrum of EVOO but not in that of VCO. The band near 3006 cm^{-1} was associated to the stretching vibration of *cis* double bonds CH groups while the latter denoted the stretching vibration of carbon-carbon double bonds. The *cis* double bonds CH groups and carbon-carbon double bonds are found widely in unsaturated fatty acids such as oleic acids, linoleic acids and linolenic acids. This correlated well with initial fatty acid profiles (Table 1) which show that EVOO

contains more unsaturated double bonds than VCO. In fact, there was only approximately 5% of fatty acids in VCO are made of unsaturated double bonds compared to over 80% of that in EVOO.

FTIR spectra for oxidized oils

In EVOO, the band near 3006 cm^{-1} experienced a progressive decrease in intensity as well as wave number throughout the period of study (Figure 4). As discussed before, this band is corresponding to the *cis* double-bond (=CH) stretching vibration therefore its intensity is affected by the number of *cis* double bonds present. Thus, the changes observed indicate a decrease in the number of *cis* double-bond, therefore a decrease in degree of unsaturation as a consequence of oxidation process. In contrast, in VCO (Figure 5), the intensity of its band near 3006 cm^{-1} remained fairly constant as only flat lines, not bands were observed throughout the study. However, this did not indicate that no oxidation occurred in VCO. As oxidation proceeds, the concentration of hydroperoxides increases and this gave a broad band which overlapped with that of the glyceride ester carbonyl groups (Gullén and Cabo, 2004). The increase in concentration of hydroperoxides can be reflected by increase band absorbance at 3470 cm^{-1} . Anyway, FTIR cannot provide sufficient quantitative data to show oxidative stability of oils. Thus, chemical analysis such as Iodine Value, Peroxide Value, Phenolic content etc need to be carried out to further investigate thermal stability of oils.

Iodine Value

IV is a measure of the degree of unsaturation in oil. The initial IV of EVOO and VCO was 81.87 ± 0.81 and 11.09 ± 0.69 respectively (Table 3). The values are similar to that reported in literature (Lide 1996). The differences in initial IV between EVOO and




Figure 4. Changes in the region near 3006 cm^{-1} of EVOO at different stages of the storage Study. (From top to bottom: raw, storage day 0, 10, 20, 30 and 40)




Figure 5. Changes in the region near 3006 cm^{-1} of VCO at different stages of the storage Study. (From top to bottom: raw, storage day 0, 10, 20, 30 and 40)

VCO are due to the different fatty acids composition. The lower IV in VCO compared to EVOO indicates that VCO is far more saturated than EVOO. This was in

good agreement with the results obtained from GC-FID as well as FTIR spectra at the band near 3006 cm^{-1} . As revealed by statistical analysis of the data, both IVs of

EVOO and VCO were decreased significantly ($P<0.05$) after 40 days of storage to 72.13 ± 0.77 and 3.26 ± 0.11 respectively. This reduction in IVs suggests a decrease in number of double bond in the oils. The maximum reductions in IVs of both oils were achieved by the heat treatment as deep frying accelerates the oxidation in the oils (Naz *et al.* 2005). In term of percentage loss, the decrease of IV in EVOO was much smaller compared to VCO. This is because of the higher total phenolic content in EVOO. IV does not take into account the structural differences present in different fatty acids such as the nature, quantity and position of the unsaturated bonds in the chain available for oxidation (Bouaid *et al.* 2007). Therefore, IV is not considered as the best way to evaluate the oxidative stability of oil. Other tests must be present to support the results and conclusions that drawn from IV.

Total phenolic content

Phenolic compounds are the main factor rendering nutritional importance to cold-pressed oil. The changes in total phenolic content of EVOO and VCO are recorded in Table 3. The initial total phenolic content of EVOO was 12.95 mg GAE per 100 g of oil. On the other hand, VCO was found to contain 0.65 mg GAE per 100 g of oil. The total phenolic content of EVOO decreased significantly from 12.95 in fresh sample to 6.94 in heated sample (0 days) and finally to 4.12 mg GAE per 100 g in the end of the experimentation (40 days after heat treatment). There were significant changes observed in VCO as well, from initial level of 0.65 to 0.51 after heat treatment (0 day) to a final value of 0.32 mg GAE per 100 g in day 40. The reduction indicates degradation of phenolic compounds in the oil samples and is related to the decrease of oxidative stability since phenolic compounds enhance the total

antioxidant activity. In this study, the total phenolic content of fresh EVOO was approximately 20 times higher than that in fresh VCO (12.95 vs. 0.65 mg GAE per 100 g of oil). In the end of the experimentation, the concentration of phenolic compounds in EVOO was still approximately 13-fold of that in VCO (4.12 vs. 0.32 mg GAE per 100 g of oil). The higher total phenolic content in EVOO at all time throughout the study explains its' better stability than VCO as revealed by relatively lower concentration of oxidative and degradation products as well as smaller percentage loss in IV at the end of experimentation. However, the percentage loss in total phenolic content of EVOO was higher than VCO (68.19% vs. 50.77%). The better retention of phenolic compounds in VCO could be explained by involvement of other antioxidants such as tocopherols and tocotrienols in counteracting oxidation process (Rossi *et al.*, 2007), which are not necessary from the family of phenolic compounds. From Iodine Value and Total Phenolic Content results, we found that both oils are stable to certain extent. Thus, we can conclude that VCO is relatively stable as compared to EVOO.

Conclusion

Heat has significantly decreased the stability of VCO and EVOO. Results from GC showed higher stability in VCO with no significant changes in proportion of each fatty acid was observed throughout the storage. While percentage of palmitic acid and linolenic acid in EVOO showed significant decrease upon storage. The occurrence of oils oxidation was proven lower in VCO through FTIR spectra. Besides that, Iodine Value (IV) and Total Phenolic content (TPC) showed reduction of 70.60% in VCO and 11.90% in EVOO while reduction of TPC, 50.77% in VCO

and 68.19% in EVOO. This reduction indicated degradation of phenolic compounds and unsaturated fatty acids in oils after heat treatment and upon storage. Thus, in term of thermal stability, we can conclude that VCO is relatively stable as compared to EVOO and is suitable for frying.

Acknowledgements

The author is thankful for the financial support from Faculty of Applied Sciences, UCSI University. Thanks also to Mr. Neo for the technical support and guides in running GC.

References

- Andrikopoulos, N.K., Kalogeropoulos, N., Falirea, A.F. and Barbagianni, M.N. 2002. Performance of virgin olive oil and vegetable shortening during domestic deep-frying and pan-frying of potatoes. International Journal of Food Science and Technology 37: 177–190.
- Association of Official American Chemists (AOAC), 1984. Official Method of Analysis, 14th edition.
- Besbes, S., Blecker, C., Deroanne, C., Lognay, G., Drira, N.E. and Attia, H. 2005. Heating effects on some quality characteristic of date seed oil. Food Chemistry 91: 469-476.
- Boskou, D. 2003. Frying Fats, Chemical and Functional Properties of Food Lipids, New York: CRC Press.
- Bouaid, A., Martinez, M. and Aracil, J. 2007. Long storage stability of biodiesel from vegetable and used frying oils. Fuel 86: 2596-2602.
- Chung, J.S., Lee, Y.S. and Choe, E. 2006. Effects of sesame oil addition to soybean oil during frying on the lipid oxidative stability and antioxidants contents of the fried products during storage in the dark. Journal of Food Science 71: 222-226.
- Fellows, P.J. 2003. Food Processing Technology: Principles and Practice, 2nd edition, CRC Press, England.
- Gomez-Alonso, S., Fregapane, G., Salvador, M.D. and Gordon, M.H. 2003. Changes in phenolic composition and antioxidant activity of virgin olive oil during frying. Food Chemistry 51: 667-672.
- Guillen, M.D. and Cabo, N. 2004. Study of the effects of smoke flavourings on the oxidative stability of the lipids of pork adipose tissue by means of Fourier transform infrared spectroscopy. Meat Science 66: 647-657.
- Lide, D.R. 1996. Handbook of Chemistry and Physics, 76th Edition, CRC Press, Boco Raton, USA.
- Lim, Y.Y., Lim, T.T. and Tee, J.J. 2007. Antioxidant properties of several tropical fruits: A comparative study. Food Chemistry 103: 1003-1008.
- Mackay, S. 2000. Techniques and Types of Fat Used in Deep-Fat Frying: A policy statement and background paper, The Heart Foundation of New Zealand.
- Naz, S., Siddiqi, R., Sheikh, H. and Sayeed, S.A. 2005. Deterioration of olive, corn and soybean oils due to air, light, heat

- and deep-frying. Food Research International 38: 127-134.
- Nevin, K.G. and Rajamohan, T. 2004. Beneficial effects of virgin coconut oil on lipid parameters and in vitro LDL oxidation. Clinical Biochemistry 37: 830-835.
- Nevin, K.G. and Rajamohan, T. 2006. Virgin coconut oil supplemented diet increases the antioxidant status in rats. Food Chemistry 99: 260-266.
- Qian, M., 2003. Gas chromatography, Food analysis laboratory manual, New York: Kluwer academic publishers.
- Rossi, M., Alamprese, C. and Ratti, S. 2007. Tocopherols and tocotrienols as free radical-scavengers in refined vegetable oils and their stability during deep-fat frying. Food Chemistry 102: 812-817.
- Rotondi, A., Bendini, A., Cerretani, L., Mari, M., Lercker, G. and Toschi, T.G. 2004. Effect of olive ripening degree on the oxidative stability and organoleptic properties of Cv. Nostrana di Brisighella extra virgin olive oil. Journal of Agricultural and Food Chemistry 52: 3649-3654.
- Villarino, B.J., DY, L.M. and Lizada, M.C.C. 2005. Descriptive sensory evaluation of virgin coconut oil and refined, bleached and deodorized coconut oil. LWT-Food Science and Technology 40: 193–199.
- Warner, K. and Mounts, T.L. 1993. Frying stability of soybean and canola oils with modified fatty acids composition. Journal of American Oil Chemistry Society 70: 983-988.
- Xu, Y.X., Hanna, M.A. and Josiah, S.J. 2007. Hybrid hazelnut oil characteristics and its potential oleochemical application. Industrial Crops and Products 26: 69-76.