

MiniReview

The effects of food processing on biogenic amines formation

Chong, C. Y., *Abu Bakar, F., Russly, A. R., Jamilah, B. and Mahyudin, N. A.

*Faculty of Food Science and Technology,
University Putra Malaysia, Serdang 43400, Selangor, Malaysia*

Abstract: Biological amines are nitrogenous compounds that occur naturally in wide variety of food. Histamine, putrescine, cadavarine, tyramine, spermine, spermidine, tryptamine and β -phenylethylamine are the biogenic amines that are normally present in foods. Although the biogenic amines play some important physiological functions but high level of amines can cause toxicological effects. High amount of amines can be produced by bacteria during amino acids decarboxylation and have been identified as one of the important agent causing seafood intoxication. Temperature is the major factor for controlling the biogenic amines formation in food. The effects of other alternatives are also discussed including salting, packaging, irradiation, high pressure processing and the use of starter culture. A variety of techniques can be combined together to control the microbial growth and enzyme activity during processing and storage for better shelf life extension and food safety.

Keywords: Biogenic amines, histamine, food processing, food safety

Introduction

Biogenic amines (BA) are the compounds in which one, two or three hydrogen of ammonia are replaced by alkyl or aryl groups (Shalaby, 1996). Figure 1 shows the chemical structure of several major biogenic amines. Putrescine, cadaverine, spermine and spermidine have aliphatic structure whereas tyramine and phenylethylamine containing aromatic structure. Heterocyclic structures are found in histamine and tryptamine (Santos, 1996). They can also be classified into monoamines (phenylethylamine and tyramine), diamines (cadaverine and putrescine) or polyamines (spermidine and spermine) based on the number of amine groups (Spano *et al.*, 2010).




Figure 1. shows the structures of biogenic amines (Önal, 2007).

Biogenic amines are natural compounds which can be produced during the normal metabolism of living cells (ten Brink *et al.*, 1990). It is also present

in food such as fish, wine, cheese, dairy product, beer, meat and vegetable. Histamine, tryptamine, β -phenylethylamine and tyramine have important physiological roles in humans (Shalaby, 1996). During food spoilage, microorganism can produce high concentration of biogenic amines by decarboxylating the free amino acids. The concentrations of biogenic amines have been suggested as indices for bacterial contamination in food (Rezaei *et al.*, 2007).

The occurrences of biogenic amines in foods (ten Brink *et al.*, 1990; Shalaby, 1996; Santos, 1996). fish (Rawles *et al.*, 1996) and dry fermented sausages (Suzzi and Gardini, 2003) have been reviewed. There is little literature on the effects of different food processing on the production of biogenic amines. The objective of this paper is to review briefly the effects of food processing on biogenic amines formation.

Outbreak and epidemiology

Biogenic amines intoxication is always related with intake of fish belongs to the *Scorbroid* family. The consumption of high amount of biogenic amines in food can result in histamine poisoning and tyramine toxicity. Histamine poisoning is the most toxic and common form of poisoning. Histamine intoxication, is also termed Scorbroid poisoning is an important food borne disease over the world. Outbreaks are common in the United States, Canada, Japan and other countries with a high consumption of fish (Behling and Taylor, 1982).

In the United States, seafood ranked third among various foods which caused food poisoning during 1983-1992. Scorbroid poisoning is the most

*Corresponding author.
Email: fatim@food.upm.edu.my

important causative agent following the consumption of mahi-mahi, tuna, mackerel, bluefish, sardines, amberjack and abalone. The major cause of scombroid poisoning is temperature abuse (Lipp and Rose, 1997). Wallace *et al.* (1999) reviewed food borne disease data gathered by the New York State Department of Health (NYSDOH) between 1980 and 1994. Among 148 seafood-associated outbreaks, scombrototoxin was one of the most common agents, responsible for 44% sea food associated disease outbreak.

In the USA, 69 outbreaks of scombrototoxin poisoning involving 297 cases were reported to CDC's Food Disease Outbreak surveillance system between 1993 and 1997 (Olsen *et al.*, 2000). From 1998 to 2002, around 118 outbreaks of scombrototoxin poisoning involving 463 cases were reported (Lynch *et al.*, 2006). In Canada, the most common seafood-associated diseases is scombroid poisoning from improperly stored fish, but Paralytic shellfish poisoning (PSP) and ciguatera poisoning have the most serious consequences (Todd, 1997). In United Kingdom, PHLS Communicable Disease Surveillance Centre reported 1425 outbreaks of Infectious Intestinal Disease (IID) from 1992 to 1999 (Gillespie *et al.*, 2001). According to this report, 10% were related with the fish and shellfish. The consumption of spoiled tuna (47%) was the most important cause for scombrototoxic fish poisoning. It was followed by mollusks contaminated with viral pathogen (36%) and crustaceans contaminated by salmonella or viral pathogens (11%).

Histamine poisoning

Scombroid fish poisoning is related with the intake of fish from *Scombroid* family such as tuna, mackerel and bonito (McLauchlin *et al.*, 2006). However, certain non-scombroid fish are also implicated in histamine poisoning including mahi-mahi, bluefish and sardines (Taylor *et al.*, 1989). FDA (2009) identify some potential species of fish related to histamine poisoning hazards including amberjack, anchovy, bluefish, bonito, oilfish, herring, jack, jobfish, mackerel, mahi-mahi, marlin, sardine, saury, shad, trevally and tuna.

Histamine fish poisoning produce one or more of the following symptoms including reddening on the face, neck and upper chest, vomiting, sweating, nausea, abdominal cramps, headache, diarrhea, dizziness, palpitations and flushing (McLauchlin *et al.*, 2006). After ingestion of the food, the incubation periods range from a few minutes to a few hours (Becker *et al.*, 2001). Histamine fish poisoning is frequently misdiagnosed due to its typical symptoms

mimic those of allergy (Attaran and Probst, 2002).

The presence of secondary amines including putrescine and cadaverine contribute to synergistic effect and enhancing the histamine toxicity in food poisoning (Taylor, 1983). Amines are also a possible mutagenic precursor, react with nitrites forming nitrosamines which is carcinogenic (Shalaby, 1996). Healthy people are able to metabolize the dietary histamine rapidly. However, the accumulation of histamine exceeding the capacity of histamine degradation can results in histamine toxicity. Clinical symptoms are more serious in people consuming drug that can retard the enzyme that metabolizes the histamine in intestine (Stratton *et al.*, 1991). Consumption of alcohol and certain drugs such as antihypertensive, antidepressants, antihypotonics, antiarrhythmics and other drugs are able to inhibit the diamine oxidase and increasing the susceptibility of people to histamine intoxication (Maintz and Novak, 2007).

The histamine detoxification systems in human comprised of diamine oxidase (DAO) and histamine N-methyl transferase. DAO plays the major role in histamine catabolism (Hungerford, 2010). Amine oxidases catalyze the oxidative deamination of biogenic amines producing the aldehyde, ammonia and hydrogen peroxide (Longu *et al.*, 2005). Diamine oxidase (DAO) catalyze the oxidative deamination of histamine and histamine-N-methyl transferase (HNMT) catalyze the ring methylation of histamine (Maintz *et al.*, 2006). DAO has been suggested for degrading histamine extracellularly whereas histamine N-methyltransferase can only metabolize histamine intracellularly (Maintz and Novak, 2007). The treatment for histamine toxicity involves the use of antihistamines (Attaran and Probst, 2002).

Effects of various foods processing on biogenic amines formation

The control of biogenic amines formation mainly focused on the controlling the growth of biogenic amines forming bacteria. It is because the histamine is heat stable (ten Brink *et al.*, 1990; Santos, 1996; Kurt and Zorba, 2009) and is not detectable organoleptically by even trained panelists (Tapingkae *et al.*, 2010). Once formed, histamine is difficult to destroy by using methods including freezing, cooking, retorting, or smoking (Etkind *et al.*, 1987). However, there are some methods are able to degrade the histamine including the gamma irradiation (Kim *et al.*, 2004) and application of diamine oxidase bacteria to degrade the histamine (Dapkevicius *et al.*, 2000).

Storage temperature is the most important factor contributing to biogenic amines formation. The effects of temperature abuse on biogenic amines formation have been studied extensively. High amount of amines can be produced under high temperature storage reported by many authors (Du *et al.*, 2002; Rodtong *et al.*, 2005; Kim *et al.*, 2002; Wei *et al.*, 1990). 25°C was optimum for histamine production by *Morganella morgani* in the artificially contaminated muscles of mackerel, albacore, and mahi-mahi (Kim *et al.*, 2002). Histamine amount increased drastically after six hours at 25°C. Kim *et al.* (2002) detected a high level of histamine of 4610 ppm in mackerel after 24 hours of storage; 3430 ppm in albacore; and 3340 ppm in mahi-mahi. Temperature of 4°C retarded the *M. morgani* growth in all species up to 14 days.

Economou *et al.* (2007) assessed the effect of temperature abuse on histamine formation in tuna muscle stored at different temperature. For storage temperature 0–2°C, 258.3 ppm histamine was detected in fresh tuna loins abused at 30°C for 2 hours daily for 12 days. This value was higher than that of the control sample of 33.5 ppm. High histamine concentration of 1962 ppm was found in tuna stored at 6–7°C after 12 days of temperature abuse storage. Icing temperature was found to retard the histamine formation. Du *et al.* (2002) only found 18 ppm of histamine in tuna after 9 days storage at 0°C whereas 68.8 ppm and 564 ppm histamine were detected in tuna stored at 4°C and 10°C. Similarly, Rodtong *et al.* (2005) only detected 19 ppm of histamine in Indian anchovy (*Stolephorus indicus*) after 15 days at ice storage.

Salting

Roseiro *et al.* (2006) studied the higher final NaCl concentrations in dry fermented pork sausage to evaluate the effects on biogenic amines levels. The author reported that the 6% salt content reduced the total biogenic amines levels significantly compared to 3% salt content. The usual formulation for the same product ranges from 4% to 4.5%. The higher salt content showed the reductions by 83%, 43%, 28% and 98% for cadaverine, putrescine, tyramine and phenylethylamine level respectively. Different salt content could be attributed to the variation of microflora composition and leading to the differences in biogenic amines formation. In this study, 6% salt concentration had lower Enterobacteriaceae, Enterococci and total aerobic psychrotrophic counts compared to 3% salt concentration. Enterobacteriaceae is responsible for the histidine decarboxylase activity to produce histamine (Bover-Cid *et al.*, 2009). In salt-fermented soybean paste, higher salt content (12%) had the lower biogenic amines level compared

to lower salt condition (6% and 8% salt) (Kim *et al.*, 2005a). Similarly, some studies found the high salt content can control the biogenic amines formation in Feta cheese (Valsamaki *et al.*, 2000) and in meat batter (Bover-Cid *et al.*, 2009).

Rodtong *et al.* (2005) identified prolific histamine producers from spoiled Indian anchovy (*Stolephorus indicus*) as *M. morgani*, *Proteus vulgaris*, and *Enterobacter aerogenes*. At 5% NaCl, these three strains were still able to produce high histamine concentration in medium. However, all isolates did not produce histamine at ≥10% NaCl. Lakshmanan *et al.* (2002) investigated the changes of the amine forming microorganisms in salt-dried sardines (*Sardinella gibbosa*). The salt content of final products ranged from 10 to 16%. The author found that only cadaverine and putrescine were produced by the bacteria isolated and did not found histamine formers during salt-drying process. The author concluded that the growth of amine forming microflora was inhibited with over 10% NaCl. However, Kongpun (2000) observed that the histamine content increased with the increase of salt content in Spanish Mackerel and achieving the highest concentration when 13-15% salt content was recorded. The author explained that it may be due to the increase of total viable count and histamine forming bacteria count with the increase of salt content.

Packaging

Modified atmosphere packaging is a popular preservation method involving the changing of gas composition surrounding the food product and packaging with barrier film. Oxygen, nitrogen and carbon dioxide are usually used in this technique and carbon dioxide is the major gas with bacteriostatic and fungistatic properties. The inclusion of carbon dioxide may inhibit the growth and increase the lag phase of microorganism with amino acid decarboxylase.

There are some studies reporting the successful inhibition of biogenic amines using modified atmosphere packaging in fish (Ozogul *et al.*, 2002a; Özogul *et al.*, 2002b; Emborg *et al.*, 2005; Ozogul and Ozogul, 2006) and chicken meat (Balamatsia *et al.*, 2006; Patsias *et al.*, 2006). The modified atmosphere packaging has a better inhibitory effect compared to vacuum packaging (Ozogul *et al.*, 2002a; Emborg *et al.*, 2005; Ozogul and Ozogul, 2006; Alak *et al.*, 2010). Emborg *et al.* (2005) studied the effect of vacuum packaging and modified packaging on biogenic amine formation at 2°C and 10°C in tuna muscle inoculated with psychrotolerant bacteria. Histamine achieved toxic level in chilled vacuum packaging tuna steaks at 2°C. But modified packaging with 40% CO₂/60%

O₂ was reported to inhibit the histamine formation. The author suggested vacuum packaged tuna may have caused the histamine intoxication during the last decade.

Ozogul and Ozogul (2006) found the modified atmosphere packaging (60% CO₂ and 40% N₂) was most effective in retarding the production of amines in sardine compared to vacuum packaging and normal air storage. The same author, Ozogul *et al.* (2002a) showed the histamine amount in herring reached 396 ppm in air, 284 ppm in vacuum packaging and 197 ppm in modified atmosphere packaging (60% CO₂ and 40% N₂) after 16 days stored at 2°C. Similarly, modified atmosphere packaging (60% CO₂ and 40% N₂) was effective in inhibiting the production of amines in herring compared to that stored in air (Özogul *et al.*, 2002b).

However, Dalgaard *et al.* (2006) found the modified atmosphere packaging (40% CO₂ and 60% N₂) did not differ significantly from air storage in reducing the histamine production in chilled fresh garfish at 0 and 5°C. Similarly, a modified atmosphere with gas composition of 60% CO₂, 25% N₂ and 15% O₂ and gas composition of 40% CO₂, 40% N₂ and 20% O₂ were not effective in reducing the amines production in hake compared to air storage (Ruiz-Capillas and Moral, 2001). But the author showed that the higher O₂ level did retard the amines formation. Gallas *et al.* (2010) also showed that the higher oxygen (75% O₂, 25% CO₂) had significant lower biogenic amine concentration in chicken meat compared to modified atmosphere of 75% N₂ and 25% CO₂.

Chitosan was studied on the use of food edible film due to its antimicrobial properties (Jeon *et al.*, 2002). Recently, chitosan film packaging was found to have the best histamine inhibitory effect in Atlantic bonito fillet and it was followed by modified atmosphere packaging (100% CO₂), vacuum packaging and cling film packaging (Alak *et al.*, 2010). The fillet packaged with chitosan film and modified packaging had significant lower histamine concentration and enterobacteriaceae count. Enterobacteriaceae was known as a major bacteria group for histamine production.

Irradiation

Irradiation is one of the important food preservation techniques. Food irradiation involves the exposure of food to ionizing radiations such as gamma rays, high energy electrons and X-rays (Arvanitoyannis *et al.*, 2009). The ionising radiation inactivates the microorganism by damaging the nucleic acid of cells (Farkas, 2006). Besides the microbial inactivation, the food irradiation is also able to induce the

radiolytic degradation of biogenic amines. More than 50 countries have adopted irradiation (Rabie *et al.*, 2010).

Irradiation was found to reduce biogenic amines content in aqueous solutions (Kim *et al.*, 2004), in fish (Mendes *et al.*, 2000; Mendes *et al.*, 2005; Mbarki *et al.*, 2008), in Chinese Rugao ham (Wei *et al.*, 2009), in low salt-fermented soybean paste (Kim *et al.*, 2005a) and in ground beef and pork (Min *et al.*, 2007). Kim *et al.* (2004) investigated the irradiation effects on amine standards dissolved in distilled water. The author demonstrated that the radiolytic degradation of biogenic amines decreased the biogenic amine significantly in a dose-dependent manner. In this study, irradiation broke down putrescine and spermine completely at 5 kGy. 10 kGy and 15 kGy were found to breakdown spermidine and histamine respectively.

In fish, irradiation significantly retarded the production of histamine, tyramine, cadaverine and putrescine in blue jack mackerel with 1 to 3 kGy (Mendes *et al.*, 2000). Mendes *et al.* (2005) showed that histamine was only detected in the control compared to irradiated sample in horse mackerel (*Trachurus trachurus*) after 23 days of ice storage. The author reported even lower level (1 kGy) of irradiation was effective to reduce the amines contents significantly. Similarly, Mbarki *et al.* (2008) reported gamma irradiation retarded the histamine production significantly ($p \leq 0.05$) in Bonito (*Sarda sarda*) with doses ranging from 0 to 7.5 kGy during chilled storage. The author showed the decrease was correlated with the increased dose ($R^2 = 0.97$). The author suggested the irradiation dose below 4 kGy was sufficient to preserve bonito quality during chilled storage.

In ripened sausages, irradiation reduced the total biogenic amines concentrations by 40%, 47% and 68% with the dosage of with 2, 4 and 6 kGy during storage (Rabie *et al.*, 2010). In Chinese Rugao ham, γ -irradiation was also reported to reduce the volatile N-nitrosamines, and residual nitrite in dry-cured ham (Wei *et al.*, 2009). Nitrite can react with putrescine and cadaverine to produce carcinogenic nitrosamines and may also cause hemoglobinaemia (Kurt and Zorba, 2009).

However, Kim *et al.* (2003) found no significant difference in biogenic amines concentration between irradiated samples and control after irradiation. Irradiation was also reported to increase concentration of phenylethylamine in pepperoni (Kim *et al.*, 2005b), spermidine, cadaverine, tryptamine and phenylethylamine in Chinese Rugao ham (Wei *et al.*, 2009). It could be explained by the radicals

produced from irradiation may alter the protein physical chemical properties and hence increase the concentration of certain biogenic amines (Rabie *et al.*, 2010). Although the irradiation is effective in controlling biogenic amines formation but it may pose some adverse effects on the aspects of food nutrition and organoleptic properties. Mbarki *et al.* (2008) found the poly-unsaturated fatty acid was reduced significantly and irradiation induced maximum lipid oxidation rates with doses of 6 and 7.5 kGy.

High pressure processing

Recently high pressure processing has become an alternative method to preserve the food. The food is subjected to high hydrostatic pressure (usually among 100 and 1000 MPa) for shelf life extension (Bárceñas *et al.*, 2010). The application of the non-thermal technology has the advantages of maintaining sensory and nutritional properties of foods compared to traditional heat treatment. This technology is being applied in the meat (Omer *et al.*, 2010), vegetable (Colle *et al.*, 2010) and seafood processing (Li *et al.*, 2009). Microorganisms are inactivated when the present of factors causing the changes of cell structure or physiological functions (Lado and Yousef, 2002). High pressure processing inactivates microorganisms by damaging membranes, denaturing the enzymes and changing the cell morphology (Murchie *et al.*, 2005).

The different levels of pressure and treatment time in high pressure processing influence the biogenic amines content. In some cases, higher pressure treatment was not effective to retard biogenic amines formation. Paarup *et al.* (2002) studied the effects of high pressure processing (15 min at ambient temperature and stored at 4°C) on biogenic amines in vacuum-packed squid mantles. For histamine, pressure of 200 MPa and 300 MPa did not show the retarding effect, its level was higher than that of control and 150 MPa. But 400 MPa was found to retard histamine production. Paarup *et al.* (2002) reported that the tyramine levels in sample of 300 and 400 MPa were higher than that of the 150 and 200 MPa and control.

Latorre-Moratalla *et al.* (2007) reported that pressure of 200 MPa (10 min at 17°C) strongly inhibit putrescine and cadavarine production in meat batter but no inhibitory effect was found on tyramine accumulation. Another study on vacuum-packaged frankfurter conducted by Ruiz-Capillas *et al.* (2007) showing that 400 MPa (10 min at 30°C) was effective to delay the tyramine, putrescine and cadavarine formation after 62 days of chilled

storage at 2°C as compared to control lot. Novella-Rodríguez *et al.* (2002a) reported 50 MPa for 72 h produced the highest amine concentrations and three times higher tyramine concentration as compared to untreated goat cheese. By contrast, the amine content of sample treated with higher pressure for short time (400 MPa for 5 min) was similar to control lot. There was study did not found the difference on biogenic amines formation in milk between pressure treatment (500 MPa for 15 min, 20°C) and heat pressurization (Novella-Rodríguez *et al.*, 2002b). The effects of high pressure processing on biogenic amines formation need more investigation.

Starter culture

In fermented food production, a starter culture is added to the raw material to accelerate the fermentation process and to obtain the better shelf life, desirable characteristics such as texture and sensory profile (Leroy and De Vuyst, 2004). The mixture of strains of lactic acid bacteria, *staphylococci* and *micrococci* genus are usually used as commercial starter cultures (Hugas and Monfort, 1997). *Staphylococcus* or *micrococcus* spp. and lactic acid bacteria are commonly applied in fermented sausages due to their lipolytic and proteolytic properties (Gücükoğlu *et al.*, 2010). The equilibrium between amines formed and degraded amines influence the biogenic amines level in food (Gardini *et al.*, 2002). Therefore, during the fermentation process, the biogenic amines formation can be controlled by using the starter cultures that are less effective in decarboxylating the amino acids to produce biogenic amines. The other approach is to use the starter cultures with amine oxidase to degrade the biogenic amines.

Various studies were done on the biogenic amines degradation by different bacteria. Mah and Hwang (2009) found that *S. xylosus* degraded 38.0% of the histamine and 4.4% of the tyramine in a phosphate buffer. Later *S. xylosus* was used as starter culture and applied to the ripening of a salted and fermented anchovy, and decrease total biogenic amines concentration by 16.0% as compared to control. Leuschner *et al.* (1998) observed that *Micrococcus* strain showed the highest tyramine oxidase activity and *Lactobacillus plantarum* had only low efficiency in degrading histamine and tyramine *in vitro*. However, Fadda *et al.* (2001) findings disagreed with Leuschner *et al.* (1998) result and found 2 strains of *L. casei* showing the greatest tyramine oxidase activity (93 and 98% degradation) and 60 and 69% degradation for 2 strains of *L. plantarum* after 96 hours of incubation in buffer system. Dapkevicius *et al.* (2000) found 4 isolate of *L. sakei* and 1 isolate

of *L. curvatus* were able to degrade 20–56% of the histamine within 30 hour in a model systems. Recently, Zaman *et al.* (2010a) isolated the *Bacillus amyloliquefaciens* and *Staphylococcus carnosus* from fish sauce with histamine degradation activity up to 59.9% and 29.1% respectively in buffer system. These cultures were used as starter culture in fish sauce fermentation, *Staphylococcus carnosus* FS19 and *Bacillus amyloliquefaciens* FS05 was found to reduce the histamine concentration by 27.7% and 15.4% as compared to control, respectively (Zaman *et al.*, 2010b).

Besides, there are studies reported that the use of starter culture is effective in inhibiting biogenic amines formation in fermented meat sausage (Gençcelep *et al.*, 2007; Gücükoğlu *et al.*, 2010), pork sausage (Lu *et al.*, 2010; Coloretti *et al.*, 2008), carp sausages (Hu *et al.*, 2007) and fish sauce (Zaman *et al.*, 2010b). Mixed starter culture of *Lactobacillus farciminis* and *Staphylococcus saprophyticus* were inoculated in fermented sausages and significantly reduced the levels of histamine, putrescine, cadaverine and tyramine compared to *Pediococcus pentosaceus* and *Staphylococcus xylosum* (Lu *et al.*, 2010). In this study, *Lactobacillus farciminis* and *Staphylococcus saprophyticus* were found to inhibit *Staphylococcus sciuri*, *Enterococcus faecalis*, *Enterobacter aerogenes*, *Lactobacillus sakei*, *Pseudomonas* sp. and *Micrococcus luteus*. *Enterobacteriaceae*, *Clostridium*, *Lactobacillus*, *Streptococcus*, *Micrococcus*, and *Pseudomonas* species possess the amino acid decarboxylase to produce the biogenic amines (Shalaby, 1996). Many enterobacteriaceae and pseudomonas are able to produce histamine, cadaverine and putrescine. Some micrococaceae and lactic acid bacteria can produce considerable amount of putrescine, histamine and tyramine (Hu *et al.*, 2007).

The mixture of starter culture of *Lactobacillus plantarum* together with *Kocuria varians* was effective in reducing the total biogenic amines concentration compared with *Lactobacillus plantarum* alone and control in low-acid salami (Coloretti *et al.*, 2008). Two different starter cultures using mixture of *Lactobacillus sakei* and *Staphylococcus carnosus* and mixture of *Pediococcus acidilactici*, *Staphylococcus xylosum* and *Lactobacillus curvatus* were effective to reduce the amounts of putrescine, cadaverine and tyramine significantly compared to control in Turkish dry-fermented sausage (Gençcelep *et al.*, 2007). Gücükoğlu *et al.* (2010) reported that the three different starter culture using *L. sakei*, *S. xylosum*, *L. plantarum*, *S. carnosus*, and *L. curvatus* were able to reduce the biogenic amines formation compared

to control in Turkish fermented sausages. In another study of silver carp sausages, three group mixed starter cultures involving *Lactobacillus plantarum*, *Staphylococcus xylosum*, *Pediococcus pentosaceus* and *Lactobacillus casei* subsp. *Casei* were able to reduce the histamine, putrescine, cadaverine and tyramine significantly after 2 days of fermentation. In this study, the use of starter culture reduced the histamine by 90–95% in sausage compared to the control (Hu *et al.*, 2007).

During chilled and room temperature storage, Komprda *et al.* (2001) found the total biogenic amines of sausage using starter culture B (*L. sakei*, *S. carnosus*, *Pediococcus pentosaceus*) was significantly higher than of starter culture A (*L. sakei*, *S. carnosus*, *S. xylosum*) at the end of the study. However, the total biogenic amines content during ripening in dry fermented sausage did not differ significantly between these two starter cultures. Komprda *et al.* (2001) suggested the different biogenic amines formation during the storage was due to the different starter culture and microflora present in sausage.

Although the use of starter culture showed a positive results in degrading or controlling the biogenic amines formation but are not necessarily effective under real manufacturing process. In fermented products, starter culture is the most important factor in influencing biogenic amines formation (Komprda *et al.*, 2009). The other chemico-physical factors including raw material, pH, aW, ripening temperature, storage temperature, sausage diameter and additives used (Komprda *et al.*, 2009). These parameters should be investigated further to determine the optimum fermentation process and least biogenic amines formation.

Conclusion

Histamine poisoning is important food borne disease and international trade issue. It can be easily misdiagnosed and not all the incidents go reported. Foods are susceptible to contamination by biogenic amines producing microorganisms during post harvest handling. High level of biogenic amines can be avoided by using good quality raw material, applying good hygienic food handling and avoiding temperature abuse during handling, delivery and storage. More studies needed to be done on the effects of irradiation and high pressure processing on biogenic amines formation and on food quality. The combination of these preservative factors (hurdle) in influencing the microbial stability, the organoleptic and nutritional quality of foods needs more investigation.

Reference

- Alak, G., Hisar, S. A., Hisar, O. and Genççelep, H. 2010. Biogenic amines formation in Atlantic bonito (*Sarda sarda*) fillets packaged with modified atmosphere and vacuum, wrapped in chitosan and cling film at 4°C. European Food Research and Technology, article in press, 1-6.
- Arvanitoyannis, I. S., Stratakos, A. and Mente, E. 2009. Impact of irradiation on fish and seafood shelf life: a comprehensive review of applications and irradiation detection. Critical Reviews in Food Science and Nutrition 49: 68–112.
- Attaran, R. R. and Probst, F. 2002. Histamine fish poisoning: a common but frequently misdiagnosed condition. Emergency Medicine Journal 19(5): 474-475.
- Balamatsia, C. C., Paleologos, E. K., Kontominas, M. G. and Savvaidis, I. N. 2006. Correlation between microbial flora, sensory changes and biogenic amines formation in fresh chicken meat stored aerobically or under modified atmosphere packaging at 4 degrees C: possible role of biogenic amines as spoilage indicators. Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology 89(1): 9–17.
- Bárceñas, M. E., Altamirano-Fortoul, R. and Rosell, C. M. 2010. Effect of high pressure processing on wheat dough and bread characteristics. LWT - Food Science and Technology 43(1): 12-19.
- Becker, K., Southwick, K., Reardon, J., Berg, R. and MacCormack, J. N. 2001. Histamine poisoning associated with eating tuna burgers. JAMA: The Journal of the American Medical Association 285(10): 1327-1330.
- Behling, A. R. and Taylor, S. L. 1982. Bacterial histamine production as a function of temperature and time of incubation. Journal of Food Science 47(4): 1311-1317.
- Bover-Cid, S., Torriani, S., Gatto, V., Tofalo, R., Suzzi, G. and Belletti, N., 2009. Relationships between microbial population dynamics and putrescine and cadaverine accumulation during dry fermented sausage ripening. Journal of Applied Microbiology 106(4): 1397-1407.
- Colle, I., Van Buggenhout, S., Van Loey, A. and Hendrickx, M. 2010. High pressure homogenization followed by thermal processing of tomato pulp: Influence on microstructure and lycopene *in vitro* bioaccessibility. Food Research International 43(8): 2193-2200.
- Coloretti, F., Chiavari, C., Armaforte, E., Carri, S. and Castagnetti, G. B. 2008. Combined use of starter cultures and preservatives to control production of biogenic amines and improve sensorial profile in low-acid salami. Journal of Agricultural and Food Chemistry 56(23): 11238-11244.
- Dalgaard, P., Madsen, H. L., Samieian, N. and Emborg, J. 2006. Biogenic amine formation and microbial spoilage in chilled garfish (*Belone belone belone*) – effect of modified atmosphere packaging and previous frozen storage. Journal of Applied Microbiology 101(1): 80-95.
- Dapkevicius, M. L. N. E., Nout, M. J. R., Rombouts, F. M., Houben, J. H. and Wymenga, W. 2000. Biogenic amine formation and degradation by potential fish silage starter microorganisms. International Journal of Food Microbiology 57(1-2): 107-114.
- Du, W. X., Lin, C. M., Phu, A. T., Cornell, J., Marshall, M. and Wei, C. I. 2002. Development of biogenic amines in yellowfin tuna (*Thunnus albacares*): effect of storage and correlation with decarboxylase-positive bacterial flora. Journal of Food Science 67: 292–301.
- Economou, V., Papadopoulou, C., Brett, M. M., Frillingos, S. and Nichols, T. 2007. Changes in histamine and microbiological analyses in fresh and frozen tuna muscle during temperature abuse. Food Additives and Contaminants 24(8): 820–832.
- Emborg, J., Laursen, B. G. and Dalgaard, P. 2005. Significant histamine formation in tuna (*Thunnus albacares*) at 2°C - effect of vacuum- and modified atmosphere-packaging on psychrotolerant bacteria. International Journal of Food Microbiology 101(3): 263-279.
- Etkind, P., Wilson, M. E., Gallagher, K. and Cournoyer, J. 1987. Bluefish-associated scombroid poisoning. JAMA: The Journal of the American Medical Association 258(23): 3409-3410.
- Fadda, S., Vignolo, G. and Oliver, G. 2001. Tyramine degradation and tyramine/histamine production by lactic acid bacteria and Kocuria strains. Biotechnology Letters 23(24): 2015-2019.
- Farkas, J. 2006. Irradiation for better foods. Trends in Food Science and Technology 17: 148–152.
- Internet: Food Administration 2009. Chapter 3: Table 3-1 Potential Vertebrate Species-Related Hazards. Downloaded from <http://www.fda.gov/Food/GuidanceDocuments/Seafood/ucm091085.htm> on 2/9/2010.
- Gallas, L., Standarová, E., Steinhauserová, I., Steinhauser, L. and Vorlová, L. 2010. Formation of biogenic amines in chicken meat stored under modified atmosphere. Acta Veterinaria Brno 79: S107–S116.
- Gardini, F., Martuscelli, M., Crudele, M. A., Paparella, A. and Suzzi, G. 2002. Use of *Staphylococcus xylosus* as a starter culture in dried sausages: effect on the biogenic amine content. Meat Science 61(3): 275-283.
- Genççelep, H., Kaban, G. and Kaya, M. 2007. Effects of starter cultures and nitrite levels on formation of biogenic amines in sucuk. Meat Science 77(3): 424-430.
- Gillespie, I.A., Adak, G.K., O'Brien, S.J., Brett, M.M. and Bolton, F.J. 2001. General outbreaks of infectious intestinal disease associated with fish and shellfish, England and Wales, 1992-1999. Communicable Disease and Public Health 4(2): 117-123.
- Güçükoğlu, A. and Küplülü, Ö. 2010. The effect of different starter cultures and ripening temperatures on formation of biogenic amine in Turkish fermented sausages. European Food Research and Technology 230(6): 875-884.

- Hu, Y., Xia, W. and Liu, X. 2007. Changes in biogenic amines in fermented silver carp sausages inoculated with mixed starter cultures. *Food Chemistry* 104(1): 188-195.
- Hugas, M. and Monfort, J. M. 1997. Bacterial starter cultures for meat fermentation. *Food Chemistry* 59(4): 547-554.
- Hungerford, J. M. 2010. Scombroid poisoning: A review. *Toxicon*, 56(2), 231-243.
- Jeon, Y. J., Kamil, J. Y. V. A. and Shahidi, F. 2002. Chitosan as an edible invisible film for quality preservation of herring and Atlantic cod. *Journal of Agriculture and Food Chemistry* 50(18): 5167-5178.
- Kim, J. H., Kim, D. H., Ahn, H. J., Park, H. J. and Byun, M. W. 2005a. Reduction of the biogenic amine contents in low salt-fermented soybean paste by gamma irradiation. *Food Control* 16(1): 43-49.
- Kim, J. H., Ahn, H. J., Lee, J. W., Park, H. J., Ryn, G. H. and Kang, L. J. 2005b. Effects of gamma irradiation on the biogenic amines in pepperoni with different packaging conditions. *Food Chemistry* 89(2): 199-205.
- Kim, J. H., Ahn, H. J., Jo, C., Park, H. J., Chung, Y. J. and Byun, M. W. 2004. Radiolysis of biogenic amines in model system by gamma irradiation. *Food Control* 15(5): 405-408.
- Kim, J. H., Ahn, H. J., Kim, D. H., Jo, C., Yook, H. S. and Park, H. J. 2003. Irradiation effects on biogenic amines in Korean fermented soybean paste during fermentation. *Journal of Food Science* 68(1): 80-84.
- Kim, S. H., Price, R. J., Morrissey, M. T., Field, K. G., Wei, C. I. and An, H. 2002. Histamine production by *Morganella morganii* in mackerel, albacore, mahi-mahi, and salmon at various storage temperatures. *Journal of Food Science* 67(4): 1522-1528.
- Kongpun, O. 2000. Histamine formation during salting of Spanish mackerel (*Scomberomoms commerson*). *Journal of Aquatic Food Product Technology* 9(1): 21-30.
- Komprda, T., Sládková, P. and Dohnal, V. 2009. Biogenic amine content in dry fermented sausages as influenced by a producer, spice mix, starter culture, sausage diameter and time of ripening. *Meat Science* 83(3): 534-542.
- Komprda, T., Neznalová, J., Standara, S. and Bover-Cid, S. 2001. Effect of starter culture and storage temperature on the content of biogenic amines in dry fermented sausage polican. *Meat Science* 59(3): 267-276.
- Kurt, S. and Zorba, O. 2009. The effects of ripening period, nitrite level and heat treatment on biogenic amine formation of "sucuk" - A Turkish dry fermented sausage. *Meat Science* 82(2): 179-184.
- Lado, B. H. and Yousef, A. E. 2002. Alternative food-preservation technologies: efficacy and mechanisms. *Microbes and Infection* 4(4): 433-440.
- Lakshmanan, R., Shakila, R. J., and Jeyasekaran, G. 2002. Changes in the halophilic amine forming bacterial flora during salt-drying of sardines (*Sardinella gibbosa*). *Food Research International* 35(6): 541-546.
- Latorre-Moratalla, M. L., Bover-Cid, S., Aymerich, T., Marcos, B., Vidal-Carou, M. C. and Garriga, M. 2007. Aminogenesis control in fermented sausages manufactured with pressurized meat batter and starter culture. *Meat Science* 75(3): 460-469.
- Leroy, F. and De Vuyst, L. 2004. Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science and Technology* 15(2): 67-78.
- Leuschner, R. G., Heidel, M. and Hammes, W. P. 1998. Histamine and tyramine degradation by food fermenting microorganisms. *International Journal of Food Microbiology* 39(1-2): 1-10.
- Li, D., Tang, Q., Wang, J., Wang, Y., Zhao, Q. and Xue, C. 2009. Effects of high-pressure processing on murine norovirus-1 in oysters (*Crassostrea gigas*) *in situ*. *Food Control* 20(11): 992-996.
- Lipp, E.K. and Rose, J.B. 1997. The role of seafood in foodborne diseases in the United States of America. *Revue Scientifique et Technique* 16(2): 620-640.
- Longu, S., Mura, A., Padiglia, A., Medda, R. and Floris, G. 2005. Mechanism-based inactivators of plant copper/quinone containing amine oxidases. *Phytochemistry* 66(15): 1751-1758.
- Lu, S., Xu, X., Zhou, G., Zhu, Z., Meng, Y. and Sun, Y. 2010. Effect of starter cultures on microbial ecosystem and biogenic amines in fermented sausage. *Food Control* 21(4): 444-449.
- Lynch, M., Painter, J., Woodruff, R. and Braden, C. 2006. Surveillance for foodborne-disease outbreaks--United States, 1998-2002. *MMWR CDC Surveillance Summaries* 55(10): 1-34.
- Mah, J. H. and Hwang, H. J. 2009. Inhibition of biogenic amine formation in a salted and fermented anchovy by *Staphylococcus xylosum* as a protective culture. *Food Control* 20(9): 796-801.
- Maintz, L. and Novak, N. 2007. Histamine and histamine intolerance. *The American Journal of Clinical Nutrition* 85(5): 1185-1196.
- Maintz, L., Bieber, T. and Novak, N. 2006. Histamine Intolerance in Clinical Practice. *Dtsch Arztebl International* 103(51-52): 3477-3483.
- Mbarki, R., Sadok, S. and Barkallah, I. 2008. Influence of Gamma Irradiation on Microbiological, Biochemical, and Textural Properties of Bonito (*Sarda sarda*) During Chilled Storage. *Food Science and Technology International* 14(4): 367-373.
- McLauchlin, J., Little, C. L., Grant, K. A. and Mithani, V. 2006. Scombrototoxic fish poisoning. *Journal of Public Health* 28(1): 61-62.
- Mendes, R., Silva, H. A., Nunes, M. L. and Empis, J. M. A. 2005. Effect of low-dose irradiation and refrigeration on the microflora, sensory characteristics and biogenic amines of Atlantic horse mackerel (*Trachurus trachurus*). *European Food Research Technology* 221(3): 329-335.
- Mendes, R., Silva, H. A., Nunes, M. L. and Empis, J. M. A. 2000. Deteriorative changes during ice storage of irradiated blue jack mackerel. *Journal of Food Biochemistry* 24(2): 89-105.
- Min, J. S., Lee, S. O., Jang, A., Jo, C. and Lee, M. 2007.

- Irradiation and organic acid treatment for microbial control and the production of biogenic amines in beef and pork. *Food Chemistry* 104(2): 791–799.
- Murchie, L. W., Cruz-Romero, M., Kerry, J. P., Linton, M., Patterson, M. F. and Smiddy, M. 2005. High pressure processing of shellfish: A review of microbiological and other quality aspects. *Innovative Food Science and Emerging Technologies* 6(3): 257-270.
- Novella-Rodríguez, S., Veciana-Nogués, M. T., Saldo, J. and Vidal-Carou, M. C. 2002a. Effects of high hydrostatic pressure treatments on biogenic amine contents in goat cheeses during ripening. *Journal of Agricultural and Food Chemistry* 50(25): 7288–7292.
- Novella-Rodríguez, S., Veciana-Nogués, M. T., Trujillo-Mesa, A. J. and Vidal-Carou, M. C. 2002b. Profile of biogenic amines in goat cheese made from pasteurized and pressurized milks. *Journal of Food Science* 67(8): 2940-2944.
- Olsen, S.J., MacKinnon, L.C., Goulding, J.S., Bean, N.H. and Slutsker, L. 2000. Surveillance for foodborne-disease outbreaks, United States, 1993-1997 MMWR CDC Surveillance Summaries 49(1): 1-62.
- Omer, M. K., Alvseike, O., Holck, A., Axelsson, L., Prieto, M. and Skjerve, E. 2010. Application of high pressure processing to reduce verotoxigenic *E. coli* in two types of dry-fermented sausage. *Meat Science* 86(4): 1005-1009.
- Önal, A. 2007. A review: Current analytical methods for the determination of biogenic amines in foods. *Food Chemistry* 103(4): 1475-1486.
- Özogul, F. and Özogul, Y. 2006. Biogenic amine content and biogenic amine quality indices of sardines (*Sardina pilchardus*) stored in modified atmosphere packaging and vacuum packaging. *Food Chemistry* 99(3): 574–578.
- Özogul, F., Taylor, K. D. A., Quantick, P. and Özogul, Y. 2002a. Changes in Biogenic Amines in Herring Stored under Modified Atmosphere and Vacuum Pack. *Journal of Food Science* 67(7): 2497-2501.
- Özogul, F., Taylor, K. D. A., Quantick, P. and Özogul, Y. 2002b. Biogenic amines formation in Atlantic herring (*Clupea harengus*) stored under modified atmosphere packaging using a rapid HPLC method. *International Journal of Food Science and Technology* 37(5): 515-522.
- Paarup, T., Sanchez, J. A., Peláez, C. and Moral, A. 2002. Sensory, chemical and bacteriological changes in vacuum-packed pressurized squid mantle (*Todaropsis eblanae*) stored at 4°C. *International Journal of Food Microbiology* 74(1-2): 1–12.
- Patsias, A., Chouliara, I., Paleologos, E. K., Savvaidis, I. and Kontominas, M. G. 2006. Relation of biogenic amines to microbial and sensory changes of precooked chicken meat stored aerobically and under modified atmosphere packaging at 4°C. *European Food Research and Technology* 223(5): 683–689.
- Rabie, M. A., Siliha, H., El-Saidy, S., El-Badawy, A. A. and Malcata, F. X. 2010. Effects of γ -irradiation upon biogenic amine formation in Egyptian ripened sausages during storage. *Innovative Food Science and Emerging Technologies* 11(4): 661–665.
- Rawles, D. D., Flick, G. J. and Martin, R. E. and Steve, L. T. 1996. Biogenic amines in fish and shellfish. *Advances in Food and Nutrition Research* 39: 329-365.
- Rezaei, M., Jafari, H., Sahari, M. A., Hosseini, H., Montazeri, N., Parviz M. and Nazarinia, A. 2007. Relation of biogenic amines and bacterial changes in ice-stored southern caspian kutum (*Rutilus frisii kutum*). *Journal of Food Biochemistry* 31(4): 541–550.
- Rodtong, S., Nawong, S. and Yongsawatdigul, J. 2005. Histamine accumulation and histamine-forming bacteria in Indian anchovy (*Stolephorus indicus*). *Food Microbiology* 22(5): 475-482.
- Roseiro, C., Santos, C., Sol, M., Silva, L. and Fernandes, I. 2006. Prevalence of biogenic amines during ripening of a traditional dry fermented pork sausage and its relation to the amount of sodium chloride added. *Meat Science* 74(3): 557-563.
- Ruiz-Capillas, C., Carballo, J. and Jiménez-Colmenero, F. 2007. Consequences of high-pressure processing of vacuum-packaged frankfurters on the formation of polyamines: Effect of chilled storage. *Food Chemistry* 104(1): 202–208.
- Ruiz-Capillas, C. and Moral, A. 2001. Effect of controlled atmosphere enriched with O₂ in formation of biogenic amines in chilled hake (*Merluccius merluccius* L). *European Food Research Technology* 212(5): 546–550.
- Santos, M. H. S. 1996. Biogenic amines: their importance in foods. *International Journal of Food Microbiology* 29(2-3): 213-231.
- Shalaby, A.R. 1996. Significance of biogenic amines to food safety and human health. *Food Research International* 29(7): 675–690.
- Spano, G., Russo, P., Lonvaud-Funel, A., Lucas, P., Alexandre, H. and Grandvalet, C. 2010. Biogenic amines in fermented foods. *European Journal of Clinical Nutrition* 64(S3): S95-S100.
- Stratton, J. E., Hutkins, R. W. and Taylor, S. L. 1991. Biogenic amines in cheese and other fermented foods. A review. *Journal of Food Protection* 54: 460–470.
- Suzzi, G. and Gardini, F. (2003). Biogenic amines in dry fermented sausages: a review. *International Journal of Food Microbiology* 88(1): 41-54.
- Tapingkae, W., Tanasupawat, S., Parkin, K. L., Benjakul, S. and Visessanguan, W. 2010. Degradation of histamine by extremely halophilic archaea isolated from high salt-fermented fishery products. *Enzyme and Microbial Technology* 46(2): 92-99.
- Taylor, S. L., Stratton, J. E. and Nordlee, J. A. 1989. Histamine poisoning (scombroid fish poisoning): an allergy-like intoxication. *Clinical Toxicology* 27 (4-5) : 225 – 240.
- Taylor, S. L. and Speckhard, M. W. 1983. Isolation of histamine-producing bacteria from frozen tuna. *Marine Fisheries Review* 45: 35–39.
- ten Brink, B., Damink, C., Joosten, H. M. L. J. and Huis in 't Veld, J. H. J. 1990. Occurrence and formation of biologically active amines in foods. *International*

- Journal of Food Microbiology 11(1): 73-84.
- Todd, E. C. D. 1997. Seafood-associated diseases and control in Canada. *Review Science Technology* 16(2): 661-672.
- Valsamaki, K., Michaelidou, A. and Polychroniadou, A. 2000. Biogenic amine production in Feta cheese. *Food Chemistry* 71(2): 259-266.
- Wallace, B. J., Guzewich, J. J., Cambridge, M., Altekruise, S. and Morse, D. L. 1999. Seafood-associated disease outbreaks in New York, 1980-1994. *American Journal of Preventive Medicine* 17(1): 48-54.
- Wei, F., Xu, X., Zhou, G., Zhao, G., Li, C. and Zhang, Y. 2009. Irradiated Chinese Rugao ham: Changes in volatile N-nitrosamine, biogenic amine and residual nitrite during ripening and post-ripening. *Meat Science* 81(3): 451-455.
- Wei, C., Chen, C. M., Koburger, J. A., Otwell, W. S. and Marshall, M. R. 1990. Bacterial growth and histamine production on vacuum packaged tuna. *Journal of Food Science* 55: 59-63.
- Zaman, M. Z., Bakar, F. A., Selamat, J. and Bakar, J. 2010a. Occurrence of biogenic amines and amines degrading bacteria in fish sauce. *Czech Journal of Food Sciences* 28(5): 440-449.
- Zaman, M. Z., Bakar, F. A., Selamat, J. and Bakar, J. 2010b. Novel starter cultures to inhibit biogenic amines accumulation during fish sauce fermentation. *International Journal of Food Microbiology*, In Press, Accepted Manuscript.