

## ***In vitro* effect on the antioxidative properties of crude extract of *Chenopodium album* in presence of the organophosphate, acephate**

\*Sanjukta, D. and Ghosh, S.

Department of Chemical Technology, University College of Science and Technology,  
University of Calcutta, 92, A.P.C. Road. Kolkata-700 009, India

**Abstract:** Acephate, an organophosphorus pesticide, is a potent free radical generator. Phenolic compounds present in leafy vegetables are recently gaining interest as good sources of natural antioxidants. The aim of this study was, thus, to evaluate the role of plant phenolics present in *Chenopodium album* as a free radical scavenger, which is a commonly consumed green leafy vegetable, against acephate toxicity in vitro, due to its free radical generation ability. An attempt was also made to identify a few of the phenolic compounds present in the crude extract through RP-HPLC. DPPH radical scavenging activity and reducing power of the leafy vegetable extract, both in the presence and absence of acephate were conducted, using concentrations of 1 µg/ml, 2.5 µg/ml and 5 µg/ml of polyphenolic compounds present in the extract and a concentration of 3 mg/ml of the pesticide. The results showed that acephate significantly reduced the radical scavenging activity of the leafy vegetable extract, in comparison to the tests containing only the vegetable extract as well as a higher concentration was required to quench the ABTS<sup>•+</sup> radicals in the presence of acephate. No significant change was observed in case of inhibition of lipid peroxidation, in a linoleic acid emulsion system, in the presence of acephate when compared to the tests containing only the plant extract. Flavonoids like rutin, quercetin and rutin hydrate were identified as few of the phenolic components present in the crude *C. album* extract. Thus, from this study, it is evident that pesticides act as a serious free radical source, whose deleterious effects can be reduced, if not totally, but to a certain extent, by such green leafy vegetables which contain a wide array of natural antioxidants in the form of polyphenols, preventing unnecessary oxidation reactions.

**Keywords:** Acephate, plant polyphenols, free radicals, radical scavenging activity, lipid peroxidation

---

### **Introduction**

Green leafy vegetables are rich sources of many nutrients and their beneficial role has partly been attributed to the antioxidant components present in them of which the major portion is formed by the flavonoids, isoflavones, lignans, catechins and isocatechins (Subhasree *et al.*, 2009; Gupta and Prakash, 2009). Flavonoids are a large group of polyphenolic antioxidants that occur in several fruits, vegetables, and beverages mainly as O-glycosides. They are efficient antioxidants capable of scavenging radical species (peroxyl radicals, hydroxyl radical, superoxide radicals) forming a phenoxy radical (Rice-Evans *et al.* 1995; Rice-Evans and Miller, 1996). Plants of the Lamiaceae family, notably rosemary, oregano, sage, mint and thyme, contain a range of potential antioxidants such as carnosol, rosemanol and carvacrol, which can contribute to the antioxidant potential of the diet (Lagouri and Boskou, 1996). Thus, the antioxidative role of such leafy vegetables has been well studied proving to have good radical scavenging activities and capability of inhibiting

lipid peroxidation. Investigations have proved their excellence in inhibiting lipid peroxidation even in heated oils when compared to artificial antioxidants like BHA, BHT, etc. (Shayamala *et al.*, 2005).

Pesticides are used extensively to eradicate pests for the enhancement of food production. However, their continuous use is causing severe toxicity amongst mankind. Intoxication with pesticides even hampers animal health. (Banerjee *et al.*, 1999; Tuzmen *et al.*, 2007; Tuzmen *et al.*, 2008). Organophosphorus pesticides have outnumbered the organochlorates in terms of their usage mainly due to their relatively low toxicity and persistence in mammalian systems (Joshi *et al.*, 2009). Long-term exposures to these pesticides cause neuronal disorders primarily through the inhibition of acetylcholine esterase (AChE) and also their lipophilicity affects the lipid bilayer thus interfering with the normal functioning of the membranes (Costa, 2006; Possamai *et al.*, 2007). Such interactions have been mainly attributed to oxidative stress caused by these organophosphates (Kovasic, 2003).

---

\*Corresponding author.  
Email: sanj\_9231@yahoo.co.in

Acephate (O, S-dimethyl acetylphosphoramidothioate), an organophosphorus insecticide, is used against various insects like thrips, jassids, aphids, ants and wasps that infect paddy, wheat, etc. Free radicals are generated under normal conditions in the mitochondrial ETC and by enzymes such as lipoxygenases, peroxidases and dehydrogenases (Halliwell and Gutteridge, 1999). However, their excessive production can cause an imbalance between the prooxidant and antioxidant status of the body cells and tissues thereby causing tissue injury, DNA damage and enzyme inactivation (Poovala *et al.*, 1998) Further, sulphur-containing organophosphates, to which acephate belongs, are S-oxidized to highly reactive intermediates within cells and tissues, which may even lead to mutagenic changes in the body cells and tissues (Farak *et al.*, 2000).

*In vivo* studies on how antioxidants in plants can protect against pesticide toxicity, have been elucidated quite vividly till date. Their role in altering the body's defensive mechanisms has been highlighted in several studies (Khan *et al.*, 2005; Kamboj *et al.*, 2006; Kaur and Sandhir, 2006; Rai and Sharma, 2007; Datta *et al.*, 2010). However, very few data is available on the activity of plant antioxidants *in vitro* in presence of pesticides. Hence, in this present study, an attempt has been made to evaluate the scavenging role of the antioxidative components present in the leafy vegetable *Chenopodium album* against the toxic effects of the organophosphate insecticide acephate, *in vitro*, based on DPPH and ABTS<sup>+</sup> radical scavenging activities, reducing activity and lipid peroxidation.

## Materials and Methods

### Chemicals

Methanol, ethanol, potassium persulfate, Folin-ciocalteau's phenol reagent, gallic acid, linoleic acid were purchased from E. Merck, India (P) Limited; Potassium ferricyanide, ethylidene diamine tetra acetic acid di-sodium salt (EDTA), ammonium thiocyanate, ferrous chloride were purchased from Sisco Research Laboratory (SRL), Mumbai, India.; Flavonoid Standards of Rutin hydrate, Rutin, Quercetin, Kaemferol, 1,1-Diphenyl-2-picrylhydrazyl stable radical and ABTS<sup>+</sup> were purchased from Sigma Chemical Co. (St. Louis, MO); Acephate (Technical Grade-85% purity) was obtained from M/S Hindustan Insecticides Limited, Delhi, India.

### Preparation of plant extract

The leafy vegetable, *Chenopodium album*, was collected from the local market and was authenticated by the Department of Botany, University of Calcutta,

Kolkata, India. The polyphenolic compounds were then extracted by following the methods developed in our laboratory (Mallick *et al.*, 2009). Briefly, the whole plants (stem and leaves) of *C. album* were washed thoroughly, oven dried and ground to powder. It was then extracted using 80:20 methanol: water. The methanolic extract was used for the various *in vitro* assays. The concentrations were calculated on the basis of the total polyphenol content. Various concentrations were used as described in each test.

### Preparation of pesticide solution

A 10 ml solution of acephate was prepared by dissolving 30 mg of the pesticide in 10 ml of double distilled water.

### Determination of polyphenols

Total polyphenol content of the leafy vegetable was determined by the Folin-Ciocalteau method as described by Matthaas (2002).

### HPLC analysis of the plant extract

The hydrolyzed sample for HPLC detection of the plant polyphenols, was prepared according to the method of Hertog *et al.* (1993). 0.5 g of the dried plant material was taken in a flask along with BHA (2g/L) and sonicated for 5 minutes in 40 ml of 65% methanol. 10 ml of 6 N HCl was added and nitrogen purged for 60 seconds. The mixture was then heated on a water bath at 90°C for 2 hrs stirring constantly. It was then cooled, filtered and sonicated for 5 minutes and then used for HPLC analysis. The components were then determined by HPLC according to Siddhuraju and Becker (2003). In brief, a gradient elution was employed for flavonoids other than catechins with a mobile phase consisting of 50 mM H<sub>3</sub>PO<sub>4</sub>, pH 2.5 (solution A), and acetonitrile (solution B) which is as follows: isocratic elution 95% A/5% B, 0-5 min; linear gradient from 95% A/5% B to 50% A/50% B, 5-55 min; isocratic elution 50% A/50% B, 55-65 min; linear gradient from 50% A/50% B to 95% A/5% B, 65-67 min; post-time 6 min before next injection. The system, WATERS 2487 is equipped with a C-18 column (Nova-Pak C18, 3.9 × 150 mm). 280 nm, 340 nm and 370 nm wavelengths were selected for the detection. The flow rate of the mobile phase was 0.8 mL/min, and the injection volume was 20 µL. The peaks were identified in comparison with authentic standards.

### DPPH stable radical scavenging activity

Different concentrations (1 µg/ml, 2.5 µg/ml and 5 µg/ml) of plant extract were taken in test tubes, volume adjusted to 1 ml by using distilled water. 5 ml of 0.1-0.2 mM methanolic solution of DPPH was added to

the test tubes and shaken vigorously. The tubes were allowed to stand at room temperature for half an hour. Another set of test tubes were also analyzed where acephate was added at a concentration of 3 mg/ml (1 ml) along with the vegetable extract. Changes in the absorbance of the sample were measured at 517 nm (Oktay *et al.*, 2003). Radical scavenging activity was expressed as the inhibition percentage and calculated using the following formula.

$$\% \text{ radical scavenging activity} = \frac{[(\text{Control OD} - \text{Sample OD}) \times 100]}{\text{Control OD}}$$

#### *ABTS<sup>+</sup> radical cation scavenging activity*

The activity was determined according to Re *et al.* (1999). 5 ml 7 mM ABTS was reacted with 88  $\mu$ l 140 mM potassium persulphate overnight in dark. Prior to use it was diluted with 50% ethanol for an initial absorbance of  $\approx$  0.7 at 734 nm. 10  $\mu$ L of vegetable extract was added to 1 ml of diluted ABTS and the change in absorbance was recorded for 5 mins at 1 sec interval. The antioxidant capacity was expressed as EC<sub>50</sub>, concentrations necessary for 50% reduction of ABTS. Similar test was conducted in presence of acephate also (1 ml of acephate was added at a concentration of 3 mg/ml to the reaction mixture).

#### *Reducing activity*

Different concentrations of extracts were mixed with 0.2M (pH 6.6) phosphate buffer (2.5 ml) and 1% potassium ferricyanide (2.5 ml). The mixture was incubated at 50 °C for 20 mins. 2.5 ml of 10% TCA was added to the mixture. 2.5 ml of this solution was mixed with distilled water (2.5ml) and 0.1% FeCl<sub>3</sub> (0.5ml) and the absorbance measured at 700 nm. Increased absorbance of the reaction mixture indicates increased reducing power. Similar test was conducted in presence of acephate also (1 ml of acephate was added to the tests at a concentration of 3 mg/ml). All analysis was run in triplicates and averaged (Yen and Chen, 1995).

#### *Inhibition of lipid peroxidation in a linoleic acid emulsion*

It was determined by the method of Takao *et al.* (1994). A reaction mixture containing 50  $\mu$ g/ml of extract, 2.3 ml of phosphate buffer (pH 7.4) and 2.5 ml of linoleic acid emulsion was incubated in the dark at 37°C. 0.1 ml of the mixture was added to 0.1 ml of 30% ammonium thiocyanate and 0.1 ml of ferrous chloride and mixed thoroughly. The absorbance was read at 500 nm after every 24 hrs. Another similar set was also kept under the same conditions, where 1ml of the pesticide at a concentration of 3 mg/ml was also added along with the plant extract.

#### *Statistical analysis*

The data was expressed as Mean  $\pm$  SD. Differences among the experimental groups were analyzed using one-way ANOVA and the comparisons between the means were carried out using the Tukey test;  $p < 0.05$  was considered as statistically significant in all the experiments.

#### **Result**

##### *Total polyphenol content and components*

In the present study, the crude extract of *Chenopodium album* was used and the active components present in it were analyzed by HPLC (Figure 1). It was observed that the crude extract mainly contained flavonoids like rutin, rutin hydrate and quercetin as major components along with some unidentified materials.

##### *DPPH stable radical scavenging activity*

The DPPH radical scavenging activity showed a significant decrease ( $p < 0.05$ ) in the presence of acephate at all three concentrations of 1  $\mu$ g/ml, 2.5  $\mu$ g/ml and 5  $\mu$ g/ml of polyphenolic compounds when compared to those without pesticide treatment (Figure 2).

##### *ABTS<sup>+</sup> radical cation scavenging activity*


A significantly high EC<sub>50</sub> value, i.e. the concentration required to quench 50% of the ABTS<sup>+</sup> free radicals, was observed in case of the acephate treated tests (12.69  $\pm$  0.021) when compared to the control without any pesticide treatment (2.15  $\pm$  0.004).

##### *Reducing activity*


The reducing activity of the plant extract showed an increase in the presence of the pesticide at the concentration of 1  $\mu$ g/ml. However, at higher concentrations of 2.5  $\mu$ g/ml and 5  $\mu$ g/ml, it showed a reduction in the activity, with a sharp significant reduction ( $p < 0.05$ ) at the highest concentration (Figure 3).

##### *Inhibition of lipid peroxidation in a Linoleic acid emulsion*

No significant change was observed in the peroxidation levels in the tests containing acephate when compared to the tests containing only the plant extract. However, the tests treated with the plant extract alone, did show a significant decrease ( $p < 0.05$ ) in lipid peroxidation rates when compared to the control containing no plant extract (Figure 4).




**Figure 1.** HPLC chromatogram of *Chenopodium album* at (A) 280 nm, (B) at 340 nm and (C) 370 nm showing 1. rutin hydrate, 2. rutin, 3 and 4. quercetin




**Figure 2.** DPPH radical scavenging activity of *C. album* extract at concentrations of 1 µg/ml, 2.5 µg/ml and 5 µg/ml both in the presence and absence of the pesticide, acephate. The values are expressed as Mean  $\pm$  SD, n=3.

\*Significantly different from the sample without pesticide treatment, at respective concentrations,  $p < 0.05$ .



**Figure 3.** Reducing activity of *C. album* extract at concentrations of 1 µg/ml, 2.5 µg/ml and 5 µg/ml both in the presence and absence of the pesticide, acephate. The values are expressed as Mean  $\pm$  SD, n=3. \* Significantly different from the sample without pesticide treatment, at respective concentrations,  $p < 0.05$ .



**Figure 4.** Inhibition of lipid peroxidation in a linoleic acid emulsion by *C. album* extract in the presence of the pesticide, acephate taking water as the control. The values are expressed as Mean  $\pm$  SD, n=3

## Discussion

In the recent years, the use of pesticides and their residual effects on the animal world has gained much importance. Pesticides have been used since ages, indiscriminately, in various forms for better food production where the role of organophosphates is vast due to their low soil persistence. Acephate, belonging to this group of organophosphates, has been implicated as a potent generator of free radicals, their basic mechanism of action, causing oxidative stress and damage to tissues (Rao *et al.*, 1991) Oxidation, or the loss of an electron, can sometimes produce these reactive substances known as free radicals that cause oxidative stress in the body. Oxidative stress has been implicated in the pathology of many diseases, inflammatory conditions, cancer and ageing.

Plant sources such as fruits, vegetables, and whole grains have been proved to impart great health benefits. These observations have led to continuous research in the field of identifying specific bioactive components in food, such as antioxidants, which may be responsible for improving and maintaining health (Packer *et al.*, 2001; Fassina *et al.*, 2004). Various forms of antioxidants such as vitamins, minerals, carotenoids, and polyphenols have been reported to be present in food stuffs (Lau *et al.*, 2006), of which, an important and powerful food antioxidant is quercetin, which is a flavonoid and more specifically a flavonol. Another flavonoid, rutin, is recently gaining much interest in the field of research and has been identified as a good antidiabetic factor (Kamalakkannan *et al.*, 2006; Kamalakkannan and Stanely, 2006).

Antioxidants are capable of stabilizing or quenching such free radicals before they can react and cause harm or can even minimize the effect, if not nullify it totally. Because oxidation is a naturally occurring process within the body, a balance with antioxidants must exist to maintain health (Rietveld

and Wiseman, 2003; Liu, 2004; Goodman *et al.*, 2004) and hence the development of several enzymatic and non-enzymatic defense mechanisms like superoxide dismutase, catalase, glutathione, glutathione reductase and so on. Antioxidants may offer resistance to oxidative stress primarily through free radical scavenging and inhibiting lipid peroxidation, thus preventing the onset of deadly diseases. Apart from the antioxidants synthesized naturally, the body requires a supplement of dietary antioxidants, which can be obtained only by the consumption of an antioxidant-rich diet. The importance of green leafy vegetables, most of which are underexploited, are proven sources of essential nutrients like vitamin A and vitamin C, carotenoids, flavonoids, phenols, etc.

DPPH assay is one of the most widely used methods for screening the antioxidant activity of plant extracts. The assay is based on the evaluation of the ability of the antioxidants to scavenge the stable radical. DPPH is a stable nitrogen-centred free radical, which produces violet colour in methanol/ethanol solution. The radicals react with suitable reducing agents, during which the electrons become paired off and the solution loses colour stoichiometrically depending on the number of electrons taken up. This loss of colour or absorbance is measured spectrophotometrically at 517 nm (Blois, 1958). In the present tests, the solution is progressively reduced to the yellow coloured product, diphenylpicryl hydrazine, with the addition of the extracts in a concentration-dependent manner. The results obtained clearly indicate the potential of the plant extracts in scavenging free radicals at high concentrations in the absence of the pesticide, acephate. However, it was also evident that acephate significantly reduced the scavenging activity of the *C. album* extract at all three concentrations possibly because of the production of the free radicals at a tremendous rate which the plant extract was unable to quench.

ABTS assay is based on the measurement of the absorbance produced by the inhibition of the ABTS<sup>+</sup> radical, which has a characteristic long wavelength absorption spectrum (Sanchez-Moreno, 2002). The results obtained clearly imply that the methanol extracts of *C. album*, inhibit the radical or scavenge the radical. However, the increase in the required concentration of the plant extract or the increased EC<sub>50</sub> value in the presence of acephate proved that the antioxidant requirement in the system needs to be enhanced to deal with the large quantities of free radicals that are being generated by the pesticide.

Lipid peroxidation is the oxidative degradation of polyunsaturated fatty acids (PUFA) involving free radicals and is a serious threat to lipid membrane

structures. Lipid peroxidation by the ammonium thiocyanate method is based on the measurement of the red complex, ferric thiocyanate at 500 nm using linoleic acid as the substrate. In the present study, free radicals that were present in the system, caused peroxidation of the lipid, which was minimized to a certain extent, almost by 50%, by the plant extract. However, the presence of acephate did not show much variation in the measurements. This was probably because the pesticides themselves had abstracted protons or interacted with free radicals rather than polyunsaturated fatty acids (PUFA), thereby protecting the latter against peroxidation and hence the antioxidative role of *C. album* could not be adjudged properly in this case.

It is thus to be concluded, that green leafy vegetables has an enormous scope as a natural antioxidant source. Their tremendous potential as dietary therapeutic agents needs extensive documentation. The deleterious effects of pesticides, a major prooxidant source, in different forms and formulations are increasing in an alarming rate and have been identified with serious health hazards. Hence, inclusion of dietary polyphenols in the form of green leafy vegetables, containing flavonoids like rutin, quercetin, etc. can avert various contemporary health challenges.

## References

- Banerjee, B. D., Seth, V., Bhattacharya, A., Pasha, S. T. and Chakraborty, A. K. 1999. Biochemical effects of some pesticides on lipid peroxidation and free radical scavengers. *Toxicological Letters* 107: 33-47.
- Blois, M. S. 1958. Antioxidant determination by the use of stable free radicals. *Nature* 26: 1199-1200.
- Costa, L. G. 2006. Current issues in organophosphate toxicology. *Clinica Chimica Acta* 366: 1-13.
- Datta, D., Dhar, P., Mukherjee, A. and Ghosh, S. 2010. Influence of polyphenolic extracts from *Enhydra fluctuans* on oxidative stress induced by acephate in rats. *Food and Chemical Toxicology* 48: 2766-2771.
- Farag, A.T., Eweidah, M.H. and El-Okazy, A.M. 2000. Reproductive toxicology of acephate in male mice. *Reproductive Toxicology* 14(5): 457-462.
- Fassina, G., Vene, R., Morini, M., Minghelli, S., Benelli, R., Noonan, D. M. and Albini, A. 2004. Mechanisms of inhibition of tumor angiogenesis and vascular tumor growth by epigallocatechin-3-gallate. *Clinical Cancer Research* 10: 4865-4873.
- Goodman, G. E., Thornquist, M. D., Balmes, J., Cullen, M. R., Meyskens, F. L., Omenn, G. S., Valanis, B. and Williams, J. H. 2004. The Beta-Carotene and Retinol Efficacy Trial: incidence of lung cancer and cardiovascular disease mortality during 6-year follow-up after stopping beta-carotene and retinol supplements. *Journal of National Cancer Institute* 96:

- 1743-1750.
- Gupta, S. and Prakash J. 2009. Studies on Indian green leafy vegetables for their antioxidant activity. *Plant Foods and Human Nutrition* 64: 39-45.
- Halliwell, B. and Gutteridge, J. M. C. 1999. Free radicals, other reactive species and disease. In: Halliwell B, Gutteridge JMC, (eds). *Free Radicals in Biology and Medicine*. 3<sup>rd</sup> ed. Oxford: Clarendon Press; pp 617-783.
- Hertog, M. G. L., Hollman, P. C. H. and van de Putte, B. 1993. Content of potentially anticarcinogenic flavonoids of tea infusions, wines, and fruit juices. *Journal of Agricultural and Food Chemistry* 41: 1242-1246.
- Joshi, A. K. R. and Rajini, P. S. 2009. Reversible hyperglycemia in rats following acute exposure to acephate and organophosphorus insecticide; Role of gluconeogenesis. *Toxicology* 257: 40-45.
- Kamalakkannan, N. and Prince, P. S. 2006. Antihyperglycaemic and antioxidant effect of rutin, apolyphenolic flavonoid, in streptozotocin- induced diabetic rats. *Pharmacology and Toxicology* 98: 97-103.
- Kamalakkannan, N. and Stanely Mainzen Prince, P. 2006. Rutin improves the antioxidant status in streptozotocin-induced diabetic rat tissues. *Molecular and Cellular Biochemistry* 293: 211-219.
- Kamboj, A., Kiran, R. and Sandhir, R. 2006. N-Acetylcysteine Ameliorates Carbofuran induced Alterations in Lipid Composition and Activity of Membrane Bound Enzymes. *Molecular and Cellular Biochemistry* 286: 107-114.
- Kaur, M. and Sandhir, R. 2006. Comparative effect of acute and chronic carbofuran exposure on oxidative stress and drug metabolizing enzymes in liver. *Drug and Chemical Toxicology* 29, 415-421.
- Khan, S. M., Sobti, R. C. and Kataria, L. 2005. Pesticide-induced alteration in mice hepato-oxidative status and protective role of black tea extract. *Clinica Chimica Acta* 358: 131-138.
- Kovacic, P. 2003. Mechanism of organophosphates (nerve gases and pesticides) and antidotes: electron transfer and oxidative stress. *Current Medical Chemistry* 10: 2705-2709.
- Lagouri, V. and Boskou, D. 1996. Nutrient antioxidants in oregano. *International Journal of Food Science and Nutrition* 47: 493-497.
- Lau, F. C., Shukitt-Hale, B. and Joseph, J. A. 2006. Beneficial effects of berry fruit polyphenols on neuronal and behavioral aging. *Journal of the Science of Food and Agriculture* 86: 2251-2255.
- Liu, R. H. 2004. Potential Synergy of Phytochemicals in Cancer Prevention: Mechanism of Action. *Journal of Nutrition* 134: 3479S-3485S.
- Mallick, B., Dhar, P. and Ghosh, S. 2009. *In vitro* antioxidative property of polyphenols present in two common aquatic leafy vegetables. *J. Indian Chem. Soc* 86: 202-204.
- Matthaus, B. 2002. Antioxidant activity of extracts obtained from residues from different oilseeds. *Journal of Agriculture and Food Chemistry* 50: 3444-3452.
- Oktay, M., Culcin, I. and Kurevioglu, O. I. 2003. Determination of *in vitro* antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *Lebensmittel-Wissenschaft und-Technologie* 36: 263-271.
- Packer, L., Weber, S. U. and Rimbach, G. 2001. Molecular aspects of  $\alpha$ -tocotrienol antioxidant action and cell signaling. *Journal of Nutrition* 131: 369S-373S.
- Poovala, V.S., Kanji, V.K., Tachikawa, H. and Salahudeen, A.K. 1998. Role of oxidant stress and antioxidant protection in acephate-induced renal tubular cytotoxicity. *Toxicology Science* 46(2): 403-409.
- Possamai, F. P., Fortunato, J. J., Feier, G., Agostinho, F. R., Quevedo, J., Filho, D. W. and Dal-Pizzol, F. 2007. Oxidative stress after acute sub-chronic malathion intoxication in Wistar rats. *Environmental and Toxicological Pharmacology* 23: 198-204.
- Rai, D. K. and Sharma, B., 2007. Carbofuran-induced oxidative stress in mammalian brain. *Molecular Biotechnology* 37: 66-71.
- Rao, P.V., Prasad, S.J. and Rajendra, W. 1991. Alterations in glycolytic and oxidative potentials of rat brain during acute and chronic acephate treatments. *Biochemistry International* 23(6): 1097-1105.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M. and Rice-Evans, C. 1999. Antioxidant activity applying ABTS radical cation decolorizing assay. *Free Radical and Biological Medicine* 26: 1231-1237.
- Rice-Evans, C., Miller, N. J., Bolwell, P. G., Bramley, P. M. and Pridham, J. B. 1995. The relative antioxidant activity of plant-derived polyphenolic flavonoids. *Free Radical Research* 22: 375-383.
- Rice-Evans, C. and Miller, N. J. 1996. Antioxidant activities of flavonoids as bioactive components of food. *Biochemical Society Transactions* 24: 790-795.
- Rietveld, A. and Wiseman S. 2003. Antioxidant effects of tea: evidence from human clinical trials. *Journal of Nutrition* 133: 3285S-3292S.
- Sanchez-Moreno, C. 2002. Methods used to evaluate the free radical-scavenging activity in foods and biological system. *Food Science and Technology International* 8(3): 121-137.
- Shayamala, B. N., Gupta, S., Lakshmi, A. J. and Prakash, J. 2005. Leafy vegetable extracts-antioxidant activity and effect on storage stability of heated oils. *Innovative Food Science Emerging Technology* 6: 239-245.
- Siddhuraju, P. and Becker, K. 2003. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera Lam*). *Journal of Agriculture and Food Chemistry* 15: 2144-2155.
- Subhasree, D., Baskar, R., Laxmi Keerthana, R., Lijina Susan, R. and Rajasekaran P. 2009. Evaluation of antioxidant potential in selected green leafy vegetables. *Food Chemistry* 115: 1213-1220.
- Takao, T., Kitatani, F., Watanabi, N., Yagi, A. and Sakata, K. 1994. A simple screening method for antioxidants and isolation of several antioxidants produced by marine bacteria from fish and shellfish. *Bioscience*

Biotechnology and Biochemistry 58: 1780-1783.

- Tuzmen, M. N., Candan, N. and Kaya, E. 2007. The evaluation of altered antioxidative defence mechanism and acetylcholinesterase activity in rat brain exposed to chlorpyrifos, deltamethrin, and their combination. *Cell Biochemistry and Function* 17: 535-540.
- Tuzmen, N., Candan, N., Kaya, E. and Demiryas N. 2008. Biochemical effects of chlorpyrifos and deltamethrin on altered antioxidative defence mechanisms and lipid peroxidation in rat liver. *Cell Biochemistry and Function* 26: 119-124.
- Yen, G. C. and Chen, H. Y. 1995. Antioxidant activity of various tea extracts in relation to their antimutagenicity. *Journal of Agriculture and Food Chemistry* 43: 27-32.