

Effect of thermal and ultraviolet treatments on the stability of antioxidant compounds in single strength pineapple juice throughout refrigerated storage

Goh, S.G., *Noranizan, M., Leong, C.M., Sew C.C. and Sobhi, B.

Department of Food Technology, Faculty of Food Science and Technology, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor Darul Ehsan, Malaysia

Abstract: Thermal treatment is commonly applied in juice manufacturing as a method to pasteurize juices. However the heat may deteriorate some of the essential compounds in the juice, especially heat-sensitive antioxidants. Therefore non-thermal treatment such as ultraviolet (UV) ray has been proposed as an alternative for pasteurization. The objective of this study was to compare the effect of thermal and UV treatments on the content of antioxidants (phenolic acids, flavonoids, carotenoids, ascorbic acids) and antioxidant capacity of single strength pineapple juice. The antioxidants stability of juices throughout 14 days of refrigerated storage was also studied. Ultraviolet treatment shows higher ascorbic acid content after treatment as compared to thermally treated single strength pineapple juice. Storage time affected the studied antioxidants, where UV treatment provided better stability to ascorbic acid content while thermal treatment provided better stability to flavonoids and carotenoids.

Keywords: Thermal, ultraviolet, pineapple juice, antioxidant, storage stability

Introduction

Pineapple (*Ananas comosus* L.) is one of the most appreciated tropical fruits around the world and is consumed as fresh fruits, juice and others. Pineapple was reported to contain polyphenolic compounds and also possess antioxidant activity (Hossain and Rahman, 2011). Studies show that compounds such as phenolics and flavonoids are responsible for providing antioxidant activity (Kahkonen *et al.*, 1999; Alothman *et al.*, 2009a; Mhatre *et al.*, 2009; Danino *et al.*, 2009). Juice extracted from pineapple can be processed into many forms such as concentrated juice, mixed juice and single strength juice. Single strength pineapple juice is categorized as juice that possess 12.5% total soluble solids, pH of 3.6, 0.54% of titratable acidity and 5.6% of suspended insoluble solid (Hongvaleerat *et al.*, 2008).

Thermal and non-thermal processes are usually applied to ensure the safety of processed juice. Thermal treatment is the most common treatment used in the food industry to ensure safety of the products. The absence of heat in non-thermal technology i.e. UV irradiation produces better quality products compared to thermal treatment (Noranizan and Benchamaporn, 2007). Irradiation is a non-thermal preservation method which involves the exposure of

food materials or products to radiation, for example ultraviolet rays. Ultraviolet light at 254 nm is lethal to most microorganisms (Bintsis *et al.*, 2000) such as *Pseudomonas* spp. on fresh meat (Stermer *et al.*, 1987) and *Salmonella typhimurium* on fish (Huang and Toledo, 1982), and it also can be applied to prolong shelf life of food such as orange juice (Tran and Farid, 2004). Ultraviolet was reported to result in 90.99% inactivation of total viable bacteria and total inactivation of yeast and mold in pineapple juice (Noranizan *et al.*, 2011).

Given the great benefits provided by antioxidants, processed fruit juices that retained high amount of antioxidants from the fresh fruits are in demand. However, some of the antioxidants are destroyed by thermal treatment (Lessin *et al.*, 1997; Lee and Coates, 2003; Cortés *et al.*, 2006; Aguilar-Rosas *et al.*, 2007; Gama and Sylos, 2007) and UV treatment (Pan *et al.*, 2004; Koutchma, 2009). The losses of antioxidants may lead to colourless products, such as reported by Chen *et al.* (1995) in carrot juice, and thus causes lower quality products. Hence, the objective of this study was to compare the effect of thermal and UV treatments on the antioxidant stability of single strength pineapple juice throughout 14 days of refrigerated storage.

*Corresponding author.

Email: noraadzahan@food.upm.edu.my

Materials and Methods

Pineapple juice production

Pineapple (*Ananas comosus*) fruits of Morris variety were purchased from Pasar Borong Selangor, Malaysia. The pineapples were washed using tap water, shells were removed and the flesh was cut into small pieces using Food Slicer (EMURA ECA-201, Japan). The pineapple juice was extracted using a milling machine (Super mass colloider, Masuko Sangyo, Japan).

Juice clarification

Extracted pineapple juice was clarified by treatment with Pectinex Ultra SP-L enzyme (0.025%, v/v) at 30°C using water bath (circulating) for 30 mins. The clarified juice was then filtered using muslin cloth and the collected juice was subjected to either thermal or UV treatment immediately.

Juice treatments

Ultraviolet treatment was performed using CiderSure 3500-B (Macedon, New York) with average power of 2.3 mW/cm², exposure duration of 3.3013 seconds and dosage applied was 7.5 mJ/cm². A preliminary study has shown that pectinase was inactivated at this UV dose. Thermal pasteurization was performed by heating the pineapple juice at 97°C for 5 mins, and this inactivated the pectinase enzyme (Sairi, 2005). Freshly clarified pineapple juice was used as the control. All treated and control samples were stored at 4±1°C before analysis.

Antioxidants extraction

Extraction was performed according to Alothman *et al.* (2009a) with slight modifications on the centrifugation speed. Antioxidants of treated pineapple juice were extracted with ethanol (70%) for 3 hrs at ambient temperature using Snaker Bath (Hotech Instrument, Taiwan) with stirring. Then, it was centrifuged at 1500 x g using Kubota 5800 Refrigerated Centrifuge (Tokyo, Japan) for 15 mins at 26°C. The supernatant was concentrated using rotary evaporator (Tokyo, Japan) at 50°C until no solvent can be collected and evaporation was continued for another 5 mins to ensure total evaporation of solvent. The concentrated sample was store at 4±1°C until analysis.

Determination of total phenolics content

Total phenolic content (TPC) analysis was performed using Folin-Ciocalteu method according to Alothman *et al.* (2009a). Wavelength used was 765nm and TPC of the sample was expressed as mg of gallic acid equivalent (GAE) / 100g sample on

fresh weight basis.

Determination of total flavonoids content

Total flavonoids content (TFC) analysis was performed according to Alothman *et al.* (2009a). Wavelength used was 510 nm and TFC of the sample was expressed as catechin equivalent (CEQ)/ 100g sample on fresh weight basis.

Determination of ascorbic acid content

Ascorbic acid content (AAC) of juice were determined according to method of AOAC Method 967.21, 45.1.14(1) using 2, 6-dichloroindophenol (AOAC, 2000).

Determination of total carotenoids content

Total carotenoids content (TCC) analysis was performed according to Moore (2003). The wavelength used was 450 nm and the TCC was calculated according to the following formula and using extinction coefficient of 2500 (Lima *et al.*, 2005):

$$\text{Total carotenoid content (mg/L)} = \frac{\text{Absorbance} \times \text{extraction volume}}{\text{Sample weight} \times 100 \times \text{extinction coefficient}}$$

Determination of antioxidant capacity

The antioxidant capacity of the extracted samples were determined using 1,1-diphenyl-2-picrylhydrazyl (DPPH) according to method of Alothman *et al.* (2009a) with modifications on the DPPH concentration. Methanolic DPPH solution (0.15 mM) was added to the concentrated sample and absorbance at 515 nm against a blank solution (methanol solution) was measured using spectrophotometer. Methanolic DPPH solution without concentrated sample was used as absorbance control. Antioxidant capacity of sample was expressed as the percentage of inhibition of DPPH (Azizah *et al.*, 2009).

Storage stability of antioxidants

Control and treated samples (thermal and UV treated) were refrigerated at 4±1°C for 14 days. Antioxidants content of these samples were analysed every 2 days. The storage duration of 14 days were chosen based on the shelf life of juices studied by Tran and Farid (2004), Baxter *et al.* (2005) and Majid *et al.* (2008).

Statistical analysis

The data obtained were analyzed and interpreted by analysis of variance (ANOVA) using MINITAB v.14 Statistical Package (Minitab Inc., State College, Pennsylvania). Values expressed were mean ±

standard deviations. Significance level was set at $p \leq 0.05$. All analysis was carried out in triplicates.

Results and Discussions

Effect of treatments on antioxidants

According to Table 1, the ascorbic acid content (AAC) between juices studied has significant differences ($p \leq 0.05$) with the sequence: thermally treated juice < UV treated juice < control. This shows that UV treatment is a better processing method to retain the heat-sensitive ascorbic acid than thermal treatment due to absence of heat in UV treatment. Significant ($p \leq 0.05$) reduction of AAC of thermally treated juices in this study was in agreement with several authors (Achinewhu and Hart, 1994; Iversen, 1999; Tiwari *et al.*, 2009; Alothman *et al.*, 2009b). Ultraviolet treatment has also been reported to reduce AAC in juices (Tran and Farid, 2004; Alothman *et al.*, 2009b; Bhat *et al.*, 2011).

As for total carotenoids content (TCC), control sample had significantly ($p \leq 0.05$) higher content than both thermal and UV treated juice. The lower TCC in thermal and UV treated juice can be explained by the thermo labile and light sensitive nature of carotenoids. The TCC of UV treated juice was slightly higher than thermally treated juice but the difference was not significant ($p > 0.05$). This might be due to the nature of double bonds in carotenoids which easily absorbed UV and then undergo the process of UV photolysis. Significant reduction of TCC in orange juice after thermal treatment was also reported by other authors (Lee and Coates, 2003; Gama and Sylos, 2007).

The total phenolic content (TPC) of both thermal and UV treated samples shows reduction but not significant as compared to control sample ($p > 0.05$). The TPC between thermal and UV treated samples have no significant differences ($p > 0.05$). In significant reduction of TPC has also been reported in thermally and UV treated pomegranate juice (Pala and Toklucu, 2011).

For flavonoids, the total flavonoids content (TFC) of thermally treated juice has no significant differences ($p > 0.05$) compared to control sample. However, slight increment in TFC was observed. Flavonoids increment has been reported where thermal treatment releases more flavonoids out of cell matrix, which might be the reason for increment of TFC of thermally treated juice in this study (Bruijn *et al.*, 2008). The TFC of UV treated juice in this study recorded insignificant changes ($p > 0.05$). However in the literature, it was reported that TFC increased after UV treatment such as in starfruit juice (Bhat *et al.*, 2011) and citrus fruit (Arcas *et al.*,

Table 1. Antioxidants content and activity of pineapple juices

	Thermally pasteurized juice	UV irradiated juice	Control (no treatment)
Ascorbic acid (mg/100g)	10.065 ± 0.798 ^b	12.673 ± 1.113 ^c	16.424 ± 0.971 ^a
Carotenoids (µg/L)	5.084 ± 0.994 ^b	5.810 ± 0.866 ^b	6.694 ± 0.334 ^a
Phenolic acids (mg GAE/100g)	41.810 ± 14.590 ^a	43.020 ± 4.380 ^a	47.130 ± 10.860 ^a
Flavonoids (mg CEF/100g)	24.933 ± 6.890 ^a	20.021 ± 5.181 ^a	22.423 ± 7.787 ^a
Antioxidant capacity (%)	19.807 ± 6.523 ^b	29.793 ± 4.991 ^{ab}	35.443 ± 4.433 ^a

Mean values having different superscript low case letter within the row are significant different ($p < 0.05$)

2000). The increment of TFC is due to the increased biosynthesis of phenylalanine ammonia lyase (PAL). After irradiation, PAL content increases (Hadwiger and Schwochau, 1971; González-Aguilar *et al.*, 2007; Charles *et al.*, 2008; Pombo *et al.*, 2011;) and this in turn increases the biosynthesis of flavonoids (González-Aguilar, 2007; Alothman *et al.*, 2009b). The reason that different result obtained in this study as compared to those in the literature was not determined, but the possible reason might be that UV dosage applied in this study was lower than authors discussed above.

The antioxidant capacity of UV treated juice was higher than thermally treated juice but with no significant difference ($p > 0.05$). There was only ascorbic acid in UV treated juice that was significantly higher ($p \leq 0.05$) than thermally treated juice.

Storage stability of antioxidants

In this study, all samples showed similar decreasing trend of TPC throughout storage but both thermal and UV treated samples had significantly ($p \leq 0.05$) higher reduction after 2 weeks storage compared to control sample (Figure 1). The TPC between thermal and UV treated juices had no significant ($p > 0.05$) difference after 2 weeks of storage. This shows that UV and thermal treatments reduced TPC after 2 weeks of storage and neither UV nor thermal treatment offered good stability to phenolic acids during storage. Reduction of TPC in thermally treated grapefruit during storage has been reported (Igual *et al.*, 2011).

The TFC between all samples studied had significant ($p \leq 0.05$) differences after 2 weeks of storage. Both thermally and UV treated samples exhibited decreasing trend during storage with thermally treated juice retaining higher flavonoids (Figure 2). The greater reduction of TCC in UV-treated juice might be due to the oxidation enzyme which cannot be inactivated by UV treatment, while thermal treatment inactivates enzymes. The reduction of TFC after storage was reported by other authors (Awad and Jager, 2000; Caro *et al.*, 2004). Flavonoids of pineapple juice in this study were unstable as it was




Figure 1. Storage stability of total phenolic content in treated and control samples throughout 14 days of refrigerated storage




Figure 2. Storage stability of totalflavonoids content in treated and control samples throughout 14 days of refrigerated storage




Figure 3. Storage stability of totalcarotenoids content in treated and control samples throughout 14 days of refrigerated storage

observed that TFC in all samples studied reduced drastically from day 2 to day 6.

The TCC between all samples studied had no significant ($p > 0.05$) difference after storage in the sequence of: UV treated juice < thermally treated juice < control sample. There was slight increment in the control sample with no significant ($p > 0.05$) difference during storage, which was also reported for kiwi fruit (Tavarini *et al.*, 2008). The TCC of thermally treated juice was relatively stable and shows no significant ($p > 0.05$) difference after storage while UV treated juice shows reduction (Figure 3). The TFC and TCC of thermally treated juice were higher than UV treated juice after 2 weeks of storage. This shows that thermal treatment offered more stability to flavonoids and carotenoids of pineapple juice in this study.

The AAC of control sample was stable over time

and had no significant ($p > 0.05$) reduction. This was in agreement with the stability of AAC in untreated orange juice during 12-day storage duration (Caro *et al.*, 2004). Throughout the whole storage period, the AAC of samples in this study followed the sequence: thermal treated juice < UV treated juice < control sample. This shows that UV treatment provides better stability to ascorbic acid than thermal treatment throughout 2 weeks of storage. Reduction of AAC has previously been reported in thermally pasteurized orange juice (Polydera *et al.*, 2003; Plaza *et al.*, 2006; Tiwari *et al.*, 2009) and pineapple juice (Achinewhu and Hart, 1994) during storage.

Conclusions

Ultraviolet treatment shows higher ascorbic acid content after treatment as compared to thermally treated single strength pineapple juice. Storage time affected the studied antioxidants, where UV treatment provided better stability to ascorbic acid content while thermal treatment provided better stability to flavonoids and carotenoids. Further studies are recommended in order to know more about the effects of UV treatment and treatment parameter on the mechanism involving antioxidants stability.

Acknowledgement

The authors acknowledge the financial support by Universiti Putra Malaysia (RUGS no. 9199939)

References

- Achinewhu, S.C. and Hart, A.D. 1994. Effect of processing and storage on the ascorbic acid (vitamin C) content of some pineapple varieties grown in the Rivers State of Nigeria. Plant Foods for Human Nutrition 46: 335-337.
- Aguilar-Rosas, S.F., Ballinas-Casarrubias, M.L., Nevarez-Moorillon, G.V., Martin-Belloso, O. and Ortega-Rivas, E. 2007. Thermal and pulsed electric fields pasteurization of apple juice: Effects on physicochemical properties and flavour compounds. Journal of Food Engineering 83: 41-46.
- Alothman, M., Bhat, R. and Karim, A.A. 2009a. Antioxidant capacity and phenolic content of selected tropical fruits from Malaysia, extracted with different solvents. Food Chemistry 115: 785-788.
- Alothman, M., Bhat, R. and Karim, A.A. 2009b. UV radiation-induced changes of antioxidant capacity of fresh-cut tropical fruits. Innovative Food Science and Emerging Technologies 10: 512-516.
- AOAC, 2000. Official Methods of Analysis of AOAC International, 17th Ed., AOAC International, Gaithersburg, MD, USA, Official Method 967.21.
- Arcas, M.C., Botía, J.M., Ortúñoz, A.M. and Rio, J.A.D.

2000. UV irradiation alters the levels of flavonoids involved in the defense mechanism of *Citrus aurantium* fruit against *Penicillium digitatum*. European Journal of Plant Pathology 106: 617-622.
- Awad, M.A. and Jager, A. 2000. Flavonoid and chlorogenic acid concentration in skin of Jonagold and Elstar apples during and after regular and ultra low oxygen storage. Postharvest Biology and Technology 20: 15-24.
- Azizah, A.H., Wee, K.C., Azizah, O. and Azizah, M. 2009. Effect of boiling and stir frying on total phenolics, carotenoids and radical scavenging activity of pumpkin. International Food Research Journal 16: 45-51.
- Baxter, I.A., Easton, K., Schneebeli, K. and Whitfield, F.B. 2005. High pressure processing of Australian navel orange juice: Sensory analysis and volatile flavour profiling. Innovative Food Science and Emerging Technologies 6: 372-387.
- Bhat, R., Ameran, S., Voon, H.C., Karim, A.A. and Tze, L.M. 2011. Quality attributes of starfruit (*Averrhoa carambola* L.) juice treated with UV radiation. Food Chemistry 127: 641-644.
- Bintsis, T., Litopoulou-Tzanetaki, E. and Robinson, R.K. 2000. Existing and potential applications of UV light in the food industry – a critical review. Journal of the Science of Food and Agriculture 80: 637-645.
- Bruijn, J., Loyola, C., Aqueveque, P., Cañumir, J., Cortéz, M. and France, A. 2008. Influence of heat treatment on the antioxidant properties of GrifolaGargal hydro-alcoholic extracts. Micología Aplicada International 20: 27-34.
- Caro, A.D., Piga, A., Vacca, V. and Agabbio, M. 2004. Changes of flavonoids, vitamin C and antioxidant capacity in minimal processed citrus segments and juices during storage. Food Chemistry 84: 99-105.
- Charles, M.T., Goulet, A. and Arul, J. 2008. Physiological basis of UV-C induced resistance to *Botrytis cinerea* in tomato fruit IV. Biochemical modification of structural barriers. Postharvest Biology and Technology 47, 41-53.
- Chen, H.E., Peng, H.Y. and Chen, B.H. 1995. Changes of carotenoids, colour and vitamin A contents during processing of carrot juice. Journal of Agricultural and Food Chemistry 43: 1912-1918.
- Cortés, C., Esteve, M.J., Rodrigo, R., Torregrosa, F. and Frígola, A. 2006. Changes of colour and carotenoids content during intensity pulsed electric field treatment in orange juices. Food and Chemistry Toxicology 44: 1932-1939.
- Danino, O., Gottlieb, H.E., Grossman, S. and Bergman, M. 2009. Antioxidant activity of 1,3-dicaffeoylquinic acid isolated from Inulaviscosa. Food Research International 42(9): 1273-1280.
- Gama, J.J.T. and Sylos, C.M. 2007. Effect of thermal pasteurization and concentration on carotenoid composition of Brazilian Valencia orange juice. Food Chemistry 100: 1686-1690.
- González-Aguilar, G.A., Zavaleta-Gatica, R. and Tiznado-Hernández, M.E. 2007. Improving postharvest quality of mango Haden by UV-C treatment. Postharvest Biology and Technology 45, 108-116.
- Hadwiger, L.A. and Schwochau, M.E. 1971. UV light-induced formation of pisatin and phenylalanine ammonia lyase. Plant Physiology 47: 588-590.
- Hongveerat, C., Cabral, L.M.C., Dornier, M., Reynes, M. and Ningsanond, S. 2008. Concentration of pineapple juice by osmotic evaporation. Journal of Food Engineering 88: 548-552.
- Hossain, M.A. and Rahman, S.M.M. 2011. Total phenolics, flavonoids and antioxidant activity of tropical fruit pineapple. Food Research International 44: 672-676.
- Huang, Y.W. and Toledo, R. 1982. Effect of high and low intensity UV irradiation on surface microbiological counts and storage-life of fish. Journal of Food Science 47: 1667-1669, 1731.
- Igual, M., García-Martínez, E., Camacho, M.M. and Martínez-Navarrete, N. 2011. Changes in flavonoid content of grapefruit caused by thermal treatment and storage. Innovative Food Science and Emerging Technologies 12: 153-162.
- Iversen, C.K. 1999. Blackcurrant nectar: Effect of processing and storage on anthocyanin and ascorbic acid content. Journal of Food Science 64: 37-41.
- Kahkonen, M.P., Hopia, A.I. and Vuorela, H.J. 1999. Antioxidant activity of plant extracts containing phenolic compounds. Journal of Agricultural Chemistry 47: 3954-3962.
- Koutchma, T. 2009. Advances in UV light technology for non-thermal processing of liquid foods. Food and Bioprocess Technology 2: 138-155.
- Lee, H.S. and Coates, G.A. 2003. Effect of thermal pasteurization on Valencia orange juice colour and pigments. Food Science and Technology 36: 153-156.
- Lessin, W.J., Catignani, G.L. and Schwartz, S.J. 1997. Quantification of cis-trans isomers of provitamin A carotenoids in fresh and processed fruits and vegetables. Journal of Agricultural and Food Chemistry 45: 3728-3732.
- Lima, V.L.A.G., Mélo, E.A., Maciel, M.I., Prazeres, F.G., Musser, R.S. and Lima, D.E.S. 2005. Total phenolic and carotenoid contents in acerola genotypes harvested at three ripening stages. Food Chemistry 90: 565-568.
- Majid, F.A.A., Gani, M.A., Talib, S.Z. and Hasim, K.K. 2008. Stability of bromelain-polyphenol complex in pineapple juice. Journal of Technology 49: 27-38.
- Mhatre, M., Tilak-Jain, J., De, S. and Devasagayam, T.P.A. 2009. Evaluation of antioxidant activity of non-transformed and transformed pineapple: A comparative study. Food and Chemical Toxicology 47: 2696-2702.
- Moore, J.P. 2003. Carotenoid synthesis and retention in mango (*Mangifera indica*) fruit and puree as influenced by postharvest and processing treatments. Florida, United States: University of Florida, MSc Thesis.
- Noranizan, M. and Benchamaporn, P. 2007. Potential of non-thermal processing for food preservation in Southeast Asian countries. ASEAN Food Journal 14(3): 141-152.
- Noranizan, M., Sharizah, S., Sew, C.C. and Karim, R. 2011. Comparison between pulsed light and ultraviolet

- treatment on microflora survival and quality attributes of pineapple juice. Universiti Malaysia Terengganu 10th International Annual Symposium (UMTAS), Kuala Terengganu, Malaysia, July 11-13, 2011, p.172.
- Pala, Ç.U. and Toklucu, A.K. (2011).Effect of UV-C on anthocuanin content and other quality parameters of pomegranate juice. Journal of Food Composition and Analysis 24(6): 790-795.
- Pan, J., Vicente, A.R., Martinez, G.A., Chaves, A.R. and Civello, P.M. 2004. Combined use of UV-C irradiation and heat treatment to improve postharvest life of strawberry fruit. . Journal of the Science of Food and Agriculture 84, 1831-1838.
- Plaza, L., Sánchez-Moreno, C., Elez-Martínez, P., Ancos, B., Martín-Beloso, O. and Cano, M.P. 2006. Effect of refrigerated storage in vitamin C and antioxidant activity of orange juice processed by high-pressure or pulsed electric fields with regard to low pasteurization. European Food Research and Technology 223: 487-493.
- Polydera, A.C., Stoforos, N.G. and Taoukis, P.S. 2003. Comparative shelf life study and vitamin C loss kinetics in pasteurized and high pressure processed reconstituted orange juice. Journal of Food Engineering 60: 21-29.
- Pombo, M.A., Rosli, H.G., Martínez, G.A. and Civello, P.M. 2011. UV-C treatment affects the expression and activity of defense genes in strawberry fruit (*Fragaria x ananassa*, Duch.). Postharvest Biology and Technology 59, 94-102.
- Sairi, M. 2005. The deacidification of pineapple juice using electrodialysis with monopolar ion exchange membranes. Johor, Malaysia: UniversitiTeknologi Malaysia, MSc thesis.
- Stermer, R.A., Lasater-Smith, M. and Brasington, C.F. 1987. Ultraviolet irradiation – an effective bactericide for fresh meat. Journal of Food Protection 50: 108-111.
- Tavarini, S., Degl'Innocenti, E., Remorini, D., Massai, R. and Guidi, L. 2008. Antioxidant capacity, ascorbic acid, total phenols and carotenoids changes during harvest and after storage of Hayward kiwifruit. Food Chemistry 107, 282-288.
- Tiwari, B.K., Donnell, C.P.O., Muthukumarappan, K. and Cullen, P.J. 2009. Ascorbic acid degradation kinetics of sonicated orange juice during storage and comparison with thermally pasteurized juice. LWT-Food Science and Technology 42: 700-704.
- Tran, M.T.T. and Farid, M. 2004. UV treatment of orange juice. Innovative Food Science and Emerging Technologies 5: 495-502.