

Microbiological quality of cooked meat products sold in Kelantan, Malaysia during Ramadhan month

^{1,2}Ain Auzureen, M. Z., ²Erkihun Aklilu, ²Mohd Azam Khan, G. K., ²Ruhil Hayati, H., ²Al Sultan, I. I. and ^{3*}Soon, J. M.

¹Faculty of Agro-Based Industry, Universiti Malaysia Kelantan, Jeli Campus, 17600 Jeli, Kelantan

²Faculty of Veterinary Medicine, Universiti Malaysia Kelantan, City Campus, 16100 Pengkalan Chepa, Kelantan

³International Institute of Nutritional Sciences and Applied Food Safety Studies, School of Sport and Wellbeing, University of Central Lancashire, Preston, UK

Article history

Received: 29 August 2015

Received in revised form:

18 March 2016

Accepted: 22 March 2016

Abstract

The Ramadhan fast is a form of worship where Muslims spend the daylight hours in a complete fast. During the month of Ramadhan, a large number of people tend to purchase food from stalls or bazaars for breaking fast. It is crucial that the foods prepared and sold are handled in a clean and safe manner. Thus, this research focused on evaluating the microbiological quality of cooked meat products (beef and chicken) from food bazaars and street-vended foods in 10 districts throughout the state of Kelantan. A total of 100 samples were collected from all 10 districts in Kelantan and were tested for coliform, *Escherichia coli*, *Salmonella* spp., and *Staphylococcus* spp. Microbiological analysis of the meat samples showed unsatisfactory results where a total of 42% of the samples were found unsatisfactory for coliform. The overall prevalence for *Staphylococcus* spp. in beef and chicken were 19.6% and 12.9%. *Escherichia coli* were detected in 23.9% of beef and 12.9% of chicken. Non-compliances for *Salmonella* were found in 13% and 9.3% of beef and chicken samples. This study determined the presence of foodborne pathogen in cooked meat products and indicated the possibilities of cross contamination and lack of hygiene during food handling.

© All Rights Reserved

Keywords

Foodborne pathogen

Food handlers

Street-vended foodFRAP

Introduction

Ramadhan is the ninth month of the Islamic lunar calendar and Muslims will spend the daylight hours in a complete fast (Dodge, 2014). Fasting refers to the condition of not eating for a varying duration of time (Fasting Center International, 2002). The Ramadhan fast is a form of worship that is a part of the five pillars of Islam, and is required of every healthy adult Muslim for a complete month (between 28-30 days) for approximately 14 hours/day (Gustaviani *et al.*, 2004). Muslims will break fast when the sun sets and it is crucial that the community observing the fasting period receives an adequate and balanced meal.

Since Ramadhan is a very special occasion, a number of street vended food stalls and bazaars will operate to sell a variety of dishes. Food bazaars in Malaysia are a congregation of food stalls in an opened area. During the fasting period, the calorie intake ranged between 1300 – 1400 cal/day (Gustaviani *et al.*, 2004) compared to a normal (without fasting) daily intake of 1800-2000 cal/day. However, the surge in calories usually occurred during the breaking of fast.

People also have less time to prepare home-cooked food and due to the fasting period, people will be more lethargic and tend to purchase take away meals. Meat products represent one of the main breaking fast dishes consumed by Muslims community. Hence it is important that the foods sold at food stalls are hygienic and safe to prevent foodborne illnesses.

Foodborne illnesses, particularly food poisoning cases are on the rise, especially during the fasting month (Soon *et al.*, 2011; Soon, 2013). This may be due to the surge in consumption of take away meals and a higher number of reported cases. Inappropriate food handling practices will result in cross contamination and/or recontamination events. Pérez-Rodríguez *et al.* (2008) defined cross contamination as “a general term which refers to the transfer, direct or indirect, of bacteria or virus from a contaminated product to a non-contaminated product” and recontamination as “contamination of food after it has been submitted to an inactivation process”. Lacking in personnel hygiene among food handlers is one of the most commonly reported practices contributing to foodborne illnesses (Lues and Van Tonder, 2007).

*Corresponding author.

Email: janmei.soon@gmail.com

Tirado and Schmidt (2000) also concluded that this substantial proportion of foodborne diseases can be attributed to food preparation practices in the domestic environment. Some of the main risk factors are inappropriate storage (32%), inadequate heat treatment (26%) and cross contamination from raw to cooked foods (25%) (Smerdon *et al.*, 2001).

Street vended foods are popular among urban people as they are inexpensive, convenient and attractive (WHO, 1996). Studies from Bangladesh (Al Mamun *et al.*, 2013), China (Liu *et al.*, 2014), Korea (Cho *et al.*, 2011), Philippines (Azanza, 2005; Manguiat and Fang, 2013), Senegal (Cardinale *et al.*, 2005), South Africa (Mosupye *et al.*, 2002; Oguttu *et al.*, 2014), Taiwan (Manguiat and Fang, 2013) reported that the microbiological quality of street vended food and beverages were found unsatisfactory. In Malaysia, a number of studies in the safety of street vended foods (Haryani *et al.*, 2007) and ready-to-eat (RTE) foods (Marian *et al.*, 2012; Jamali *et al.*, 2013) had been conducted. This was followed by a few other studies on Knowledge, Attitudes and Practices (KAP) (Toh and Birchenough, 2000; Noor-Azira *et al.*, 2012; Norrakiah and Siow, 2014), food handlers' attitude at school canteens (Saidatul and Hayati, 2013) hand hygiene practices (Tan *et al.*, 2014) and food service hygiene factors (Ungku *et al.*, 2011). It is crucial that the foods prepared and sold are handled in a clean and safe manner. Thus, this research focused on evaluating the microbiological quality of cooked meat products (beef and chicken) from food bazaars and street-vended foods. So far to our knowledge the present study represents the first microbiological quality survey of cooked meat products sold in food bazaars during the Ramadhan.

Materials and Methods

Study sites and sampling

A total of 53 bazaars from all 10 districts in the state of Kelantan were selected. Kelantan has the highest Muslim population in Malaysia. The districts include: Kota Bharu, Bachok, Pasir Puteh, Tumpat, Pasir Mas, Machang, Tanah Merah, Jeli, Gua Musang and Kuala Krai (Figure 1). A total of 46 beef and 54 chicken samples were collected in July 2014 for laboratory analysis. The 100 samples were purchased from all 53 bazaars and were selected based on availability and variability of types of cooked meat products. A description of some selected cooked meat products sold at bazaars is shown in Table 1.

Laboratory procedures for meat samples

Meat samples were collected in sterile bags




Figure 1. Map showing the number of bazaars where beef and chicken meats were sampled in Kelantan, Malaysia

and transported to the laboratory in a carrier box containing ice packs. Analyses were performed upon receipt of samples at the laboratory. However, if a laboratory analysis was postponed due to delayed arrival of samples, the samples were refrigerated at 0–4°C until examination but were not kept longer than 36 hours (Al Mamun *et al.*, 2013). 25 g of each sample were homogenised in 1% buffered peptone water in a Stomacher 400 Circulator (Seward, UK) blender for 2 minutes. Following homogenization, all meat samples were tested for coliform bacteria, *Escherichia coli*, *Salmonella* spp., and *Staphylococcus* spp. Total coliform were enumerated using multiple tube fermentation technique. MacConkey and Eosin methylene blue agar were used to determine the presence of *E. coli* followed by indole tests. Rappaport Vasiliadis broth was used as selective broth for enrichment of *Salmonella* spp. and Xylose lysine deoxycholate agar was used as selective agar for detection of *Salmonella* spp. Suspected black colonies were sub-cultured to obtain pure colonies and confirmed with Triple sugar iron (TSI) agar tests. Mannitol salt agar was used as selective medium for *Staphylococcus* spp. Acid production as the result of fermentation of mannitol results in formation of yellow colonies and zones (APHA, 2001). Coliform counts of less than 10² per g were considered acceptable (ICMSF, 1986). *E. coli* should be < 3 cfu/g, coagulase positive staphylococci should be < 10² per g and *Salmonella* spp. should be absent in 25 g (EC No 2073/2005; FSANZ, 2001).

Table 1. Description of cooked meat products sold at bazaars during Ramadhan

Cooked meat products	Brief description
Black pepper beef	Beef marinated in soy sauce and black pepper
Soy sauce beef	Beef fried with soy sauce, onion and red pepper
Fried beef	Beef marinated with salt, onions and pepper and then fried
Beef / chicken <i>kurma</i>	Beef or chicken cooked in <i>kurma</i> gravy made from mixed curry powder, spices, potatoes, coconut milk and coriander.
<i>Singgang</i> beef	Beef cooked in sauce with plenty of herbs such as galangal, chillies, garlic, onion and black pepper
Spicy red beef / chicken	Beef or chicken cooked in concentrated sauce of dried chillies
Beef / chicken <i>gulai</i>	<i>Gulai</i> is similar to curry except lighter in taste and colour.
Beef / chicken <i>kerutuk</i>	Beef or chicken cooked in mixture of <i>kerisik</i> (toasted grated coconut) and <i>kerutuk</i> spices (coriander powder, cardamom seeds, clove, fennel seeds, cumin, black peppercorns, turmeric, galangal, lemongrass bulbs and garlic)
Beef <i>gulai acar</i> / <i>dalca</i>	Beef or chicken cooked in curry powder with potatoes, carrots, eggplants, green beans, chillies, curry leaves and baby corns.
Beef <i>kawah</i>	Beef cooked in a large pot of curry. Popular during wedding ceremonies in Kelantan.
<i>Airasam perut lembu</i>	Cow intestines are sliced and boiled with vinegar, lime, chives, onion, shrimp paste, chillies and tamarind
<i>Kunyit</i> beef	Beef marinated with salt and turmeric powder and fried.
Gearbox soup	Made from bull's joints and boiled in richly flavoured soup.
Ginger chicken	Chicken marinated and cooked with salt and sliced ginger.
Roasted chicken	Chicken marinated with honey, black pepper, aniseed, soy sauce, oyster sauce and ginger before roasted.
Chicken <i>Tom yam</i>	Chicken cooked in mixed spicy chilli paste with lime leaf and lemongrass. Tom yam originates from Thailand.
<i>Percik</i> chicken	Chicken cooked in coconut milk, dried chillies, garlic and lemongrass and then roasted.
Paprika chicken	Chicken cooked with lime leaves, <i>tom yam</i> paste, lemongrass, hot pepper, fish sauce, ginger, onion, garlic, sweet soy sauce and some vegetables.
<i>Kerisik</i> chicken	Chicken cooked with <i>kerisik</i> (toasted, grated coconut), galangal, chillies, ginger and brown sugar
Honey chicken butts / wings	Chicken butts / wings marinated with honey, ginger, oyster sauce, soy sauce, black pepper, and garlic and roasted.
<i>Ayam tiga rasa</i>	Chicken cooked with ginger, plum sauce, sweet soy sauce, tomato sauce, lime, spicy pepper and ginger to produce sweet, sour and spicy taste.
<i>Ayam peparu kicap</i>	Chicken lungs cooked with soy sauce, onion, garlic and capsicum.

Results and Discussion

Of the total meat samples (n=100), 62% were tested positive for total coliforms. 42% were found to be unsatisfactory (total coliforms ≥ 102 per g). Results revealed that all samples from Tanah Merah were unsatisfactory. On the other hand, all samples from Pasir Puteh and Machang districts were found satisfactory (Figure 2). Table 2 shows the overall prevalence of *Staphylococcus* spp. was 16%. There was significance difference in the prevalence among the districts (DF = 9, $p < 0.05$). The overall prevalence of *Staphylococcus* spp. in chicken was 12.9% while beef was 19.6%. The overall prevalence of unsatisfactory quality beef and chicken was 8.7%

and 3.7%. These are meat products contaminated with *Staphylococcus* spp. at concentration greater than 10^2 per g. Coagulase tests were carried out and tested negative for *S. aureus*.

In the samples tested, no coagulase positive staphylococci were detected. Coagulase positive staphylococci such as *S. aureus* cause food poisoning and superficial skin infections (Chakraborty *et al.*, 2011). However staphylococci can be routinely isolated from humans and associated environments. Staphylococci are ubiquitously distributed in man's environment and strains present in the nose often contaminate the back of hands, fingers and face (Garcia *et al.*, 1986; Lues and Van Tonder, 2007). Most food sellers did not wear gloves, masks or

Table 3. *E. coli* and *Salmonella* spp. in cooked meat samples

Districts	Isolated bacteria			
	<i>E. coli</i>		<i>Salmonella</i> spp.	
	Beef	Chicken	Beef	Chicken
Kota Bharu	62.5% (5/8)	25% (2/8)	12.5% (1/8)	37.5% (3/8)
Bachok	50% (2/4)	0% (0/4)	25% (1/4)	0% (0/4)
Jeli	50% (2/4)	25% (2/8)	0% (0/4)	12.5% (1/8)
Pasir Puteh	0% (0/5)	0% (0/5)	0% (0/5)	0% (0/5)
Gua Musang	0% (0/4)	25% (1/4)	0% (0/4)	0% (0/4)
Machang	0% (0/4)	0% (0/5)	50% (2/4)	20% (1/5)
Kuala Krai	33.3% (1/3)	0% (0/6)	33.3% (1/3)	0% (0/6)
Tanah Merah	0% (0/5)	20% (1/5)	0% (0/5)	0% (0/5)
Tumpat	25% (1/4)	25% (1/4)	0% (0/5)	0% (0/5)
Pasir Mas	0% (0/5)	0% (0/5)	20% (1/5)	0% (0/5)
Total	23.9% (11/46)	12.9% (7/54)	13% (6/46)	9.3% (5/54)

from slaughter plant (Soon *et al.*, 2011). Besides applying correct food handling techniques, on-farm intervention strategies to reduce microbial load are crucial to reduce contamination in the food chain.

Meanwhile, the presence of *Salmonella* spp. in RTE foods may be a result of undercooking, poor handling practices and cross contamination (NSW Food Authority, 2009). Cooked foods are vulnerable if touched by *Salmonella*-contaminated fingers that have been contaminated by low numbers of the bacteria (Guzewich and Ross, 1999). A person may carry *Salmonella* in their faeces without any signs of infection. They may then contaminate food by not washing their hands after using the toilet thus spreading *Salmonella* to others through contaminated food. There was no reported outbreak of typhoid fever in Kelantan during the Ramadhan period. It is possible that sporadic cases occurred but were unreported (Soon *et al.*, 2011).

Cross contamination from food contact surfaces to cooked meat products

Cross contamination via cooking utensils, food handlers, processing equipments, deficient hygiene practices, inadequate cooking and storage are closely related to foodborne outbreaks (Carrasco *et al.*, 2012). Inappropriate food handling practices such as using the same cutting board for raw and RTE food is a potential vehicle for cross contamination. *Listeria monocytogenes* (Goh *et al.*, 2014) and *Campylobacter jejuni* (Tang *et al.*, 2011) were transmitted from raw chicken meat to cooked chicken meat via cutting boards. Cutting boards are commonly perceived

as significant fomites in cross contamination of foodstuffs with foodborne agents (Carrasco *et al.*, 2012). But studies by Moore *et al.* (2007) underlined that food contact surfaces that are “easy to clean” (e.g. Formica and stainless steel) may be more likely to release foodborne pathogens during common food preparation practices.

Additionally, foodborne pathogens readily transmit from wet kitchen sponges to stainless steel surfaces to food (Kusumaningrum *et al.*, 2003) and from poultry meats to stainless steel surfaces (Malheiros *et al.*, 2010). In fact, Kusumaningrum *et al.* (2003) and Takahashi *et al.* (2011) found that pathogens remain viable on dry stainless steel surfaces and present a contamination hazard for considerable periods of time. Pests particularly flies are potential vectors for pathogens. Pest control practices observed at most bazaars include usage of candles, hand-made fly swat or adhesive paper to trap flies.

Hand washing is easy to do and it's one of the most effective ways to prevent the spread of many types of infection and illness in all setting (CDC, 2013). This is because the hands of food handlers can be vector to spread harmful microorganism through cross contamination. Food handlers can also spread microorganisms during and after they experience gastrointestinal infections (Baş *et al.*, 2006). Training is crucial to any food safety systems. Poor staff training in food hygiene is a real threat to food safety; hence effective training is an important prerequisite to successful implementation of a food safety management system (Arvanitoyannis and Kassaveti, 2009). To be effective, food safety training

needs to target changing the behaviour. Griffith (2000) argued that behavioural change (i.e. the implementation of required hygiene practices) is not easily achieved and that consideration must be given to motivation, constraints, barriers and facilities as well as to cultural aspects. Food safety practices will only be implemented given adequate resources and appropriate management culture (Clayton and Griffith, 2008). Besides educating and training in appropriate food handling practices, food handlers or operators can be trained in simple qualitative risk assessments (risk matrix: severity x probability) to determine food safety risks (Manning and Soon, 2013).

Conclusion

Food stalls and bazaars fulfil the demands of consumers and assist in socio-economic growth of food vendors. However, the safety of food sold may be compromised due to unhygienic handling and inappropriate storage temperature. Hence, priority should be placed in assisting food vendors in understanding the importance and requirements of food safety. All food sellers and handlers should be registered and trained under the Food Handlers' Training Programme and foodstalls inspected for hygiene.

Acknowledgements

The authors gratefully acknowledge the research funding from RAGS (R/RAGS/A07.00/00295A/001/2013/000120) and the Malaysia Ministry of Education for the financial support. In addition, we are grateful to the Faculty of Veterinary Medicine, Universiti Malaysia Kelantan for their support and utilization of research facilities.

References

- Al Mamun, M., Rahman, S. M. M. and Turin, T. C. 2013. Microbiological quality of selected street food items vended by school-based street food vendors in Dhaka, Bangladesh. *International Journal of Food Microbiology* 166(3): 413-418.
- APHA2001. APHA technical committee on microbiological methods for foods. *Compendium of methods for the microbiological examination of foods*, 4th Ed. APHA, Washington, D. C.
- Arvanitoyannis, I. S. and Kassaveti, A. 2009. HACCP and ISO 22000 – A comparison of the two systems. In Arvanitoyannis, I. S. (Ed.). *HACCP and ISO 22000: Application to foods of animal origin*, p. 3-45. Oxford: Wiley-Blackwell Limited.
- Azanza, P. V. 2005. Aerobic plate counts of Philippine ready-to-eat foods from take-away premises. *Journal of Food Safety* 25(2): 80-97.
- Baş, M., Ersun, A. S. and Kivanç, G. 2006. The evaluation of food hygiene knowledge, attitudes, and practices of food handlers in food businesses in Turkey. *Food Control* 17(4): 317-322.
- Cardinale, E., Perrier Gros-Claude, J. D., Tall, F., Guye, E. F. and Salvat, G. 2005. Risk factors for contamination of ready-to-eat street-vended poultry dishes in Dakar, Senegal. *International Journal of Food Microbiology* 103(2): 157-165.
- Carrasco, E., Morales-Rueda, A. and García-Gimeno, R. M. 2012. Cross-contamination and recontamination by *Salmonella* in foods: A review. *Food Research International* 45(2): 545-556.
- CDC 2013. Center for disease control and prevention. Wash your hands. Retrieved on December 12, 2014 from CDC Website: <http://www.cdc.gov/features/handwashing/>
- Chakraborty, S. P., Mahapatra, S. K. and Roy, S. 2011. Biochemical characters and antibiotic susceptibility of *Staphylococcus aureus* isolates. *Asian Pacific Journal of Tropical Biomedicine* 1(3): 212-216.
- Cho, J.-I., Cheung, C.-Y., Lee, S.-M., Ko, S.-I., Kim, K.-H., Hwang, I.-S., Kim, S.-H., Cho, S.-Y., Lim, C.-J., Lee, K.-H., Kim, K.-S. and Ha, S.-D. 2011. Assessment of microbial contamination levels of street-vended foods in Korea. *Journal of Food Safety* 31(1): 41-47.
- Clayton, D. A., and Griffith, C. J. 2008. Efficacy of an extended theory of planned behaviour model for predicting caterer's hand hygiene practices. *International Journal of Environmental Health Research* 18(2): 83-98.
- Dodge, C. H. 2009. *Understanding Islam Book. A complete guide to Muslims beliefs, practices and culture*. 2nd edn. Ohio: Adams Media.
- EC No 2073/2005 2005. Commission regulation on microbiological criteria for foodstuffs. Retrieved on November 26, 2014 from <http://fsrio.nal.usda.gov/sanitation-and-quality-standards/microbiological-standards-and-guidelines>
- Fasting Center International 2002. Fasting. Retrieved on March 20, 2015 from <http://www.hendrickhealth.org/healthy/000531.htm>
- FSANZ 2001. Guidelines for the microbiological examination of ready-to-eat foods. Food Standards Australia New Zealand. Retrieved on November 26, 2014 from <http://fsrio.nal.usda.gov/sanitation-and-quality-standards/microbiological-standards-and-guidelines>
- Garcia, M. L., Francisco, J. J. and Moreno, B. 1986. Nasal carriage of *Staphylococcus* species by food handlers. *International Journal of Food Microbiology* 3(2): 99-108.
- Goh, S. G., Leili, A.-H., Kuan, C. H., Loo, Y. Y., Lye, Y. L., Chang, W. S., Soopna, P., Najwa, M. S., Tang, J. Y. H., Yaya, R., Nishibuchi, M., Nakaguchi, Y. and Son, R. 2014. Transmission of *Listeria monocytogenes* from raw chicken meat to cooked chicken meat through cutting boards. *Food Control* 37: 51-55.

- Griffith, C. 2000. Food safety in catering establishments. In Farber J. M. and Todd, E. C. D. (Eds.). *Safe Handling of Foods*, p. 235-256. New York: Marcel Dekker, Inc.
- Gustaviani, R. R., Soewondo, P., Semiardji, G. and Sudoyo, A. W. 2004. The influence of calorie restriction during the Ramadan fast on serum fructosamine and the formation of beta hydroxybutirate in Type 2 diabetes mellitus patients. *Acta Medica Indonesiana* 36(3): 136-141.
- Guzewich, J. and Ross, M.P. 1999. Evaluation of risks related to microbiological contamination of ready-to-eat food by food preparation workers and the effectiveness of interventions to minimize those risks. Food and Drug Administration, Center for Food Safety and Applied Nutrition. Retrieved on March 24, 2015 from http://www.handwashingforlife.com/files/rte_fd_prep_risk_eval.pdf
- Haryani, Y., Noorzaleha, A. S., Fatimah, A. B., Noorjahan, B. A., Patrick, G. B., Shamsinar, A. T., Laila, R. A. S. and Son, R. 2007. Incidence of *Klebsiella pneumoniae* in street foods sold in Malaysia and their characterization by antibiotic resistance, plasmid profiling, and RAPD-PCR analysis. *Food Control* 18(7): 847-853.
- Hennekinne, J., De Buyser, M. and Dragacci, S. 2012. *Staphylococcus aureus* and its food poisoning toxins: characterization and outbreak investigation. *FEMS Microbiology Reviews* 36(4): 815-836.
- Jamali, H., Chai, L. C. and Thong, K. L. 2013. Detection and isolation of *Listeria* spp. and *Listeria monocytogenes* in ready-to-eat foods with various selective culture media. *Food Control* 32(1): 19-24.
- Hyeon, J. Y., Chung, G. T., Bing, S. H., Kwon, K. S., Lee, H. H., Kim, S. J., Jeon, S. E., Kang, Y. H. and Kim, J. 2013. A foodborne outbreak of *Staphylococcus aureus* associated with fried chicken in Republic of Korea. *Journal of Microbiology and Biotechnology* 23(1): 85-87.
- Kusumaningrum, H. D., Riboldi, G., Hazeleger, W. C. and Beumer, R. R. 2003. Survival of foodborne pathogens in stainless steel surfaces and cross contamination to foods. *International Journal Food Microbiology* 85(3): 227-236.
- Liu, Z., Zhang, G. and Zhang, X. 2014. Urban street foods in Shijiazhuang city, China: Current status, safety practices and risk mitigating strategies. *Food Control* 41: 212-218.
- Lues, J. F. R. and Van Tonder, I. 2007. The occurrence of indicator bacteria on hands and aprons of food handlers in the delicatessen sections of a retail group. *Food Control* 18(4): 326-332.
- Malheiros, P., Passos, C., Casarin, L., Serraglio, L. and Tondo, E. C. 2010. Evaluation of growth and transfer of *Staphylococcus aureus* from poultry meat to surfaces of stainless steel and polyethylene and their disinfection. *Food Control* 21(3): 298-301.
- Manguiat, L. S. and Fang, T. J. 2013. Microbiological quality of chicken- and pork-based street-vended foods from Taichung, Taiwan, and Laguna, Philippines. *Food Microbiology* 36(1): 57-62.
- Manning, L. and Soon, J. M. 2013. Mechanisms for assessing food safety risks. *British Food Journal* 115(3): 460-484.
- Marian, M. N., Sharifah Aminah, S. M., Zuraini, M. I., Son, R., Maimunah, M., Lee, H. Y., Wong, W. C. and Elexson, N. 2012. MPN-PCR detection and antimicrobial resistance of *Listeria monocytogenes* isolated from raw and ready-to-eat foods in Malaysia. *Food Control* 28(2): 309-314.
- Moore, G., Blair, I. S. and McDowell, D. A. 2007. Recovery and transfer of *Salmonella typhimurium* from four different domestic food contact surfaces. *Journal of Food Protection* 70(10): 2273-2280.
- Mosupye, F.M., Lindsay, D., Damelin, L.H. and Von Holy, A. 2002. Cytotoxicity assessment of *Bacillus* strains isolated from street-vended foods in Johannesburg, South Africa. *Journal of Food Safety* 22(2): 95-105.
- Noor-Azira, A.-M., Mohammad-Faid, A.-R., Shuhaimi, M., Syafinaz, A.-N., Rukman, A. H. and Malina, O. 2012. Knowledge, attitude and practices regarding food hygiene and sanitation of food handlers in Kuala Pilah, Malaysia. *Food Control* 27(2): 289-293.
- Norrakiah, A. S. and Siow, O. N. 2014. Knowledge, attitudes and practices of food handlers on food safety in food service operations at the Universiti Kebangsaan Malaysia. *Food Control* 37: 210-217.
- NSW Food Authority 2009. Microbiological quality guide for ready-to-eat foods. A guide to interpreting microbiological results. Retrieved on November 27, 2014 from http://www.foodauthority.nsw.gov.au/Documents/science/microbiological_quality_guide_for_RTE_food.pdf
- Oguttu, J. W., McCrindle, C. M. E., Makita, K. and Grace, D. 2014. Investigation of the food value chain of ready-to-eat chicken and the associated risk for staphylococcal food poisoning in Tshwane Metropole, South Africa. *Food Control* 45: 87-94.
- Pérez-Rodríguez, F., Castro, R., Posada-Izquierdo, G. D., Valero, A., Carrasco, E., Carcía-Gimeno, R. M. and Zurera, G. 2010. Evaluation of hygiene practices and microbiological quality of cooked meat products during slicing and handling at retail. *Meat Science* 86(2): 479-485.
- Pérez-Rodríguez, F., Valero, A., Carrasco, E., Carcía-Gimeno, R. M. and Zurera, G. 2008. Understanding and modelling bacterial transfer to foods: A review. *Trends in Food Science and Technology* 19(3): 131-144.
- Saidatul A. A. A. and Hayati, M. D. 2013. Food handlers' attitude towards safe food handling in school canteens. *Procedia - Social and Behavioral Sciences* 105: 220 - 228.
- Saif, J. M. A., Norrakiah, A. S., Aminah, A. and Ratna, D. A. R. 2009. Microbiological quality of selected ready-to-eat food at Hulu Langat district, Malaysia. *Prosiding Seminar Kimia Bersama UKM-ITB VIII*, p. 421-433. Bangi: Universiti Kebangsaan Malaysia.
- Smerdon, W. J., Adak, G. K., O'Brien, S. J., Gillespie, I. A., and Reacher, M. 2001. General outbreaks of infectious intestinal disease linked with red meat, England and Wales, 1992-1996. *Communication*

- Disease and Public Health 4(4): 259-267.
- Soon, J. M. 2013. Do more spot checks on food. The Star newspaper, 26 July, 2013.
- Soon, J. M., Chadd, S. A. and Baines, R. N. 2011. *Escherichia coli* O157:H7 in beef cattle: On farm contamination and control measures. Animal Health Research Reviews 12(2): 197-211.
- Soon, J. M., Singh, H. and Baines, R. 2011. Foodborne disease in Malaysia: A review. Food Control 22(6): 823-830.
- Takahashi, H., Kuramoto, S., Miya, S. and Kimura, B. 2011. Desiccation survival of *Listeria monocytogenes* and other potential foodborne pathogens on stainless steel surfaces is affected by different food soils. Food Control 22(3-4): 633-637.
- Tan, S. L., Bakar, F. A., Abdul Karim, M. S., Lee, H. Y. and Mahyudin, N. A. 2013. Hand hygiene knowledge, attitudes and practices among food handlers at primary schools in Hulu Langat district, Selangor (Malaysia). Food Control 34(2): 428-435.
- Tan, S. L., Lee, H. Y., Abu Bakar, F., Abdul Karim, M. S., Rukayadi, Y. and Mahyudin, N. A. 2013. Microbiological quality on food handlers' hands at primary schools in Hulu Langat District, Malaysia. International Food Research Journal 20(5): 2973-2977.
- Tan, S. L., Lee, H. Y. and Mahyudin, N. A. 2014. Antimicrobial resistance of *Escherichia coli* and *Staphylococcus aureus* isolated from food handler's hands. Food Control 44: 203-207.
- Tirado, C. and Schmidt, K. 2000. WHO surveillance programme for control of foodborne infections and intoxications in Europe, 7th report, 1993-1998. BGVVFAO/WHO Collaborating Centre for Training and Research in Food Hygiene and Zoonoses. Retrieved on December 12, 2014 from http://www.bfi.bund.de/internet/7threport/7threphome_fr.htm
- Toh, P. S. and Birchenough, A. 2000. Food safety knowledge and attitudes: culture and environment impact on hawkers in Malaysia: Knowledge and attitudes are key attributes of concern in hawker food handling practices and outbreaks of food poisoning and their prevention. Food Control 11(6): 447-452.
- Tang, J. Y. H., Nishibuchi, M., Nakaguchi, Y., Ghazali, F. M., Saleha, A. A. and Son, R. 2011. Transfer of *Campylobacter jejuni* from raw to cooked chicken via wood and plastic cutting boards. Letters in Applied Microbiology 52(6): 581-588.
- Ungku, Z. A. U. F., Huey, C. B., Murali, S. and Rosli, S. 2011. Foodservice hygiene factors – The consumer perspective. International Journal of Hospitality Management 30(1): 38-45.
- Viswanathan, P. and Kaur, R. 2000. Prevalence and growth of pathogens on salad vegetables, fruits and sprouts. International Journal of Hygiene and Environmental Health 203(3): 205-213.
- WHO. 1996. Essential requirements for street vended foods. Food Safety Unit, WHO, Geneva. Retrieved on March 24, 2014 from http://apps.who.int/iris/bitstream/10665/63265/1/WHO_FNU_FOS_96.7.pdf